Published online 2018 August 7.

**Research Article** 

# Comparative Evaluation of Conventional and Nanosilver-Containing Leucocyte and Platelet-Rich Fibrin/Biomaterial in the Anti-Biofilm Formation of Standard Species of *Candida* and *Streptococcus*

Pardis Haddadi<sup>1</sup>, Hooman Khorshidi<sup>1</sup>, Saeed Raoofi<sup>1</sup>, Ali Dehghani Nazhvani<sup>2</sup> and Parisa Badiee<sup>3,\*</sup>

<sup>1</sup>Department of Periodontology, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran <sup>2</sup>Department of Oral and Maxillofacial Pathology, Biomaterial Research Center, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran <sup>3</sup>Alborzi Clinical Microbiology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>c</sup> Corresponding author: Professor Alborzi Clinical Microbiology Research Center, Namazee Hospital, Zand Ave., 7193711351, Shiraz, Iran. Tel: +98-7136474292, Fax: +98-7136474303, Email: badieep@gmail.com

Received 2018 March 08; Revised 2018 July 29; Accepted 2018 August 01.

# Abstract

**Background:** Biomaterial products like Leukocyte and platelet-rich fibrin (L-PRF) membranes can be used in a variety of medical fields to increase the anti-infective defense system and promote tissue regeneration and wound healing due to platelet growth factors. It is one of the oldest approaches of regenerative tissue. In addition, antifungal and bacterial activities against some microorganisms are reported in the literature for silver nanoparticles.

**Objectives:** The aim of this study was to evaluate the anti-biofilm formation of L-PRF and its modification with silver nanoparticles (AgNPL-PRF) against microorganisms-associated human health conditions.

**Methods:** Blood samples were obtained and L-PRF and AgNPL-PRF membranes were produced using blood centrifugation. The biofilm formation was evaluated by incubating the standard *Candida* and *Streptococcus* species with the membranes. The incubated membranes were rinsed to remove unattached cells and incubated in an RPMI 1640 medium. The populations of growing microorganisms in the medium were evaluated by ELISA reader.

**Results:** The difference in the presence of microorganisms in RPMI medium demonstrated a statistically significant inhibition of biofilm formation of *Candida albicans* and *C. parapsilosis* while it was insignificant in *C. glabrata* and *Streptococcus mitis* in both types of membranes.

**Conclusions:** Silver nanoparticles-treated L-PRF as a biological material presented the more inhibition of biofilm formation of contaminating microorganisms and it can be used as anti-infectious material in the surgical sites.

Keywords: Candida albicans, C. parapsilosis, C. glabrata, Streptococcus mitis, Silver Nanoparticles, Leukocyte and Platelet Rich Fibrin

# 1. Background

In the modern medicine, the development of biomaterial sciences and the use of autologous materials are therapeutic approaches to enhance wound healing. One of the important biomaterials is platelet. It is a reservoir of the growth factors in its alpha-granules, which play an important role in the healing of damaged tissues. Many biomaterial products like platelet-rich plasma and leukocyteand platelet-rich fibrin (L-PRF) are the second-generation of platelet concentrate for the topical use in the treatment of soft tissue diseases and burns, hard-to-heal wounds, oral and maxillofacial surgery, and implantology (1, 2). In the fresh blood, L-PRF is prepared with leukocytes and a fibrin network after activation. In medicine, the use of plateletfibrin concentrates is one of the oldest approaches of regenerative tissue (2) and the combination of leucocytes and platelets is an interesting healing biomaterial with microbicidal activity. The cell composition of platelet-rich fibrin includes platelets equal to 97% and leukocytes  $\pm$  50% of initial blood (3).

Many surgical processes may fail due to infections and the corresponding inflammation; therefore, prevention is important in the patients' outcome. Infections may be caused by the penetration of microorganisms (bacteria or fungi) into the surgical sites depending on the type of causative agents and host immune system. *Candida* species are one of the most prevalent colonized organisms in the mouth, gastrointestinal tract, vagina, and skin, which can cause systemic infections with high morbidity in immunocompromised patients (4, 5). *Streptococcus mitis* 

Copyright © 2018, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited

is another inhabited organism in the human mouth commonly found in the throat and nasopharynx as a colonized organism causing infection in the immunocompromised patients with moderate or severe clinical diseases (6-8).

Some colonized microorganisms due to surface charge and hydrophobicity are able to make cell-cell interactions and cell surface biofilm. Biofilm formation is associated with increased production of antigens and secondary metabolites, the development of antibiotic resistance, and rates of genetic exchange, which can cause clinical problems. In maxillofacial and periodontal surgeries or during the implantation, these biofilms production act as the major risk factors for infections. The use of anti-infective biomaterials such as L-PRF in surgical sites can help the surgeons to improve the healing of the surgical wound.

Antifungal and bacterial activities against some clinical isolates and ATCC microorganisms are reported in the literature for silver nanoparticles (AgNPs) (9-11). Accordingly, the combination of fluconazole with AgNPs enhances the antifungal activity of this agent (11). However, the potential antifungal effects of L-PRF have seldom been studied.

# 2. Objectives

The aim of this study was to evaluate the antibiofilm formation of conventional and modified silver nanoparticle-containing L-PRF membranes on the standard *Candida* and *Streptococcus* species.

#### 3. Methods

## 3.1. Ethics Statement

The study protocol was in accordance with the ethical guidelines of the 1975 Helsinki declaration and approved by the ethics committee of Shiraz University of Medical Sciences (code#14866).

## 3.2. Preparing Silver Nanoparticles Suspension

Silver nanoparticles suspension was prepared by adding 200 mg of the powder (Sigma, Aldrich, USA) with a particle size of < 100 nm to 1 cc of sterile normal saline and sonicated for 2 minutes at 200 watts in a sonication device (Dr. Heilscher, UP 200H, ultraschall prozessor) (12). The serial dilutions with concentrations of 200, 100, 50, 25, 12.5, 6.25 mg/mL were prepared.

## 3.3. L-PRF and AgNPL-PRF Preparation

Venous whole blood (19 mL) was collected from 18 healthy male volunteers with the age range of 25 - 35 years. The exclusion criteria in this study were a history of any systemic diseases, smoking, and taking any medication or using anticoagulants within the last three months. For each nanosilver concentration, blood samples were divided into two tubes in sterile conditions, one tube containing 10 mL blood for the preparation of the conventional L-PRF and another containing 9 mL blood plus 1 mL silver nanoparticle suspension for the preparation of the modified L-PRF (AgNPL-PRF). Different concentrations of silver nanoparticles were provided from 20 mg/mL to 0.625 mg/mL in whole blood samples. In each concentration, three blood samples were used and L-PRF and AgNPL-PRF membranes were prepared and compared. The tubes were immediately centrifuged for 12 minutes at a low speed (2700 rpm, Eppendorf 5702 centrifuge, Hamburg, Germany), according to the protocol provided by Dohan Ehrenfest et al. (13). The fibrin membranes containing platelets were formed in the buffy coat part and middle of the tubes between the red blood cells and plasma. The fibrin membranes (Figure 1) were removed and placed on a lattice plate in PRF BOX (Process, France) and pressurized under the metal plate for 10 minutes. Each membrane was aliquoted to 4 mm in length and 2 mm in diameter and used in the study process.

#### 3.4. Evaluation of Antibiofilm Characteristics

Three Candida standard species, Candida albicans PTCC 50027 (ATCC 1023), Candida parapsilosis (ATCC 22019), Candida glabrata (ATCC 2001), and Gram-positive Streptococcus mitis (ATCC 49456) were evaluated in this study. The biofilm formations were carried based on Hassan A et al. (14). Candida species and S. mitis were cultured on Sabouraud dextrose agar (Merck, Germany) and blood agar (liofilchem, Italia), respectively. Organism suspensions with a density equal to 0.5 McFarland standards were prepared in fresh RPMI1640 (Sigma, St. Louis, Missouri). 0.5 McFarland standard was equal to 1 - 2  $\times$  10<sup>8</sup> CFU/mL for S. mitis and 1 - 2  $\times$  10<sup>6</sup> CFU/mL for Candida species. Three aliquots of different concentrations of AgNPL-PRF and L-PRF from each donor were incubated with 1 mL of 0.5 Mc-Farland suspensions of the entire tested microorganisms at 37°C. After 24 hours, supernatants were aspirated carefully and the membranes were washed twice with phosphate buffer saline (pH 7.3) in order to remove unattached cells. Only the microbial biofilm formed on the clot was evaluated. The biofilm formation of each membrane was evaluated by crystal violet (0.1%) staining (14).



Figure 1. Conventional L-PRF (A) and AgNPL-PRF (B) prepared from the human venous blood.

The quality of microorganism's biofilm formation was evaluated by adding 500 mL of RMPI 1640 medium incubated at 37°C for 24 hours. During this time, the adherent microorganisms started to multiply and scatter daughter cells in the medium. The population of the released daughter cells was evaluated, 200  $\mu$ L of each medium was poured into 96-well plates, and monitored by an ELISA Reader device (Thermolab system, Multiskan Ascent, Finland) at 490 nm. If the biofilm production of microorganisms was partially or completely inhibited, only a little growth or no growth could be observed. All membranes were tested in triplicate. RPMI medium without any microorganism and 0.5 McFarland microorganism suspensions were considered as negative and positive controls, respectively, and tested under the same conditions as mentioned. Since during the washing process, few microorganisms might be simply attached to the membranes, not to form a biofilm, we defined the index value for this study. An index value equal to one was defined as optical density (OD) of the positive control divided by that of the negative control. The index value for each concentration was calculated by dividing the mean OD of each concentration of membrane by the mean OD of the negative control. Index values were interpreted as negative (< 0.3), positive (= 1), and moderate (0.3-0.9).

# 3.5. Statistical Analysis

Data were collected in SPSS (version 15). Statistical analysis was performed using paired t-test at a significance level of P > 0.05.

# 4. Results

Crystal violet staining showed biofilms around some of the membranes. In general, the inhibition of biofilms formation was in the range of 0.1 to 1 for the L-PRF and AgNPL-PRF in different concentrations depending on the type of microorganism. After 24 hours, no inhibition of biofilm formation in C. albicans was observed in the presence of L-PRF. When using AgNPL-PRF at concentrations of 20 and 10 mg/mL, a significant decrease in biofilm formation was observed. However, no significant differentiation was seen between 10 and 20 mg/mL concentrations of silver nanoparticles in the blood samples. As shown in Figure 2, when the concentration of silver nanoparticles decreased, the growth inhibition was independent of the concentration. The index score for C. parapsilosis in L-PRF showed a moderate inhibition in biofilm formation ability while in AgNPL-PRF with concentrations of 20 and 10 mg/mL, a significant inhibition in biofilm ability was observed, compared to the lower silver nanoparticle concentrations.



Figure 2. Distribution of the anti-biofilm formation for various concentrations in different membranes. The vertical and horizontal lines represent the index value and nanosilver concentration of the membranes, respectively. There is a significant difference between the concentrations in *Candida albicans* and *C. parapsilosis* and no significant differences were found in *C. glabrata* and *Streptococcus mitis*.

L-PRF and AgNPL-PRF inhibited the biofilm formation in *S. mitis* and *C. glabrata*, independent of the silver nanoparticles concentration. For almost all the experimental conditions, the growth inhibition was similar.

## 5. Discussion

In this study, the antibacterial and antifungal activities of two biomaterials, i.e., L-PRF and AgNPL-PRF, were evaluated against biofilm formation of the standard species of *C. albicans, C. parapsilosis, C. glabrata*, and *S. mitis.* We found that in vitro antifungal activities of L-PRF and AgNPL-PRF were not similar and standard L-PRF presents a lower antifungal biofilm formation than AgNPL-PRF in *C. albicans* and *C. parapsilosis.* 

Healing the surgical wounds is the most desirable procedure in all types of surgeries like implantation in dentistry. During the healing phase, several factors are needed to inhibit the infections and its inflammation and induce cellular proliferation. Platelets are known for their role in the healing of the surgical site due to their growth factors that prevent infection due to antibacterial and fungicidal proteins stored in their granules and blood loss at the site of tissue injury (13). In the families of platelet concentrates, L-PRF contains significant concentrations of leukocytes with a stable dense fibrin matrix and platelet growth factors (15). This fibrin biomaterial represents a new opportunity to improve both the maturation of bone grafts and the final aesthetic result of the peri-implant soft tissue. Therefore, a topical application of L-PRF is used in the treatment of chronic ulcers as an inexpensive, simple, and safe agent for all recalcitrant skin ulcers (16). One of the important roles of L-PRF is the successful dental implant due to the ability to stimulate bone regeneration during dental implants (17). In many clinical trials, the efficacy of this biomaterial has proven for postoperative pain, localized site development, promotion of the soft tissue healing, sinus lift or socket preservation procedures and reduced early adverse effects of the inflammation in surgeries of oral and maxillofacial (18-20).

The use of antimicrobial agents at the site of infections to prevent infections in surgical sites may lead to increased resistant species and intolerable cytotoxic effects of hypersensitivity reactions in the patients. Health care providers should try to minimize spreading resistant etiologic agents by proper management of the antimicrobial use. To do so, the focus should be on the loading of a modified biomaterial like AgNPL-PRF. The antibacterial and antifungal activities of silver nanoparticles have been reported in the literature (21, 22) and enhanced antifungal activity against C. albicans was reported due to the combination of silver nanoparticles and fluconazole (11). The antibacterial properties of silver nanoparticles with bone cement and amalgam were reported in the literature (19, 20). The antimicrobial activity of silver nanoparticle depends on its concentration. Silver nanoparticles showed potent antifungal activity against some Candida species at concentrations of 1-7  $\mu$ g/mL (9). In the present study, the combination of L-PRF and silver nanoparticles presented similar or more effective anti-biofilm formation in the studied microorganisms compared to conventional L-PRF. The lowest concentration was 10 mg/mL of blood samples. The differentiation in concentrations may be due to the entrapped silver nanoparticles in the fibrin membrane and the sedimentation of some nanoparticles in the tube during the centrifugation of the blood samples and preparation of AgNPL-PRF.

There are populated bacteria and fungi in the human oral cavity, on which the status of oral health depends. Candida species are colonized organisms in some parts of the human body; in immunocompromised patient's postsurgery, they can pass through the natural body barriers to cause disseminated infections with high morbidity and mortality. Oral candidiasis was diagnosed in 8% of the pediatric patients, and 46.8% of the patients with hematologic disorders were colonized with Candida species (23, 24). Candida albicans is the most commonly implicated organism recovered from 60% of dentate patient's mouth over the age of 60 and 50.5% of Candida species isolated from the oral cavity of liver transplant recipients (25-27). However, other Candida species like C. parapsilosis and C. glabrata may be identified as the cause of infection (28). In this study, L-PRF did not present anti-C. albicans biofilm activity after 24 hours incubation but presented a moderate ability in *C. parapsilosis* and complete activity in *C. glabrata*. AgNPL-PRF inhibited the fungal biofilm activity on all studied Candida species, depending on silver nanoparticles concentration.

Streptococcus mitis is a member of the viridans streptococci causing some clinical diseases such as toxic shocklike syndrome, meningitis, and sepsis in immunocompromised patients. It is a normal flora of the human body, like the female genital and gastrointestinal system, oropharynx, and the skin (29, 30). According to the literature, *S. mitis* is the predominant species in soft tissue surfaces of the oral cavity and maxillary and mandibular lips (7, 8, 30). In this study, two types of the membrane presented anti-*S. mitis* biofilm formation.

## 5.1. Conclusions

As revealed, both types of L-PRF presented anti-biofilm formation depending on the type of microorganism. How-

ever, AgNPL-PRF biomaterial could inhibit the biofilm formation of all studied microbial populations and it may serve as a more suitable agent for the prevention of tissue infections compared to L-PRF. The lower concentration was 10 mg/mL of blood samples. We suggest that the use of AgNPL-PRF could help the favorable prevention of postsurgery infections in different parts of the human body. This is the first study on modified L-PRF and further research is needed to evaluate the application of this biomaterial.

## Acknowledgments

This study was supported by Prof. Alborzi Clinical Microbiology Research Center, Shiraz University of Medical Sciences. The authors would like to thank Dr. Hassan Khajea Ph.D., for copy editing of the manuscript, Hadis Jafarian and Fatemeh Ghasemi for helping with lab procedures.

# Footnotes

Authors' Contribution: Pardis Haddadi and Parisa Badiee participated in research concept/design. Pardis Haddadi, Parisa Badiee, Hooman Khorshidi, and Saeed Raoofi conducted data analysis and interpretation. Pardis Haddadi, Parisa Badiee, and Ali Dehghani Nazhvani performed the experiments and contributed to the sample processing and material synthesis. Pardis Haddadi and Parisa Badiee drafted the manuscript. Hooman Khorshidi, Saeed Raoofi, and Ali Dehghani Nazhvani conducted the critical revision of the manuscript.

**Conflict of Interests:** We declare that we have no financial interests related to the material in the manuscript.

**Funding/Support:** The funding body had no role in the study design and collection, analysis, and interpretation of data, or in writing the manuscript.

#### References

- Lubkowska A, Dolegowska B, Banfi G. Growth factor content in PRP and their applicability in medicine. J Biol Regul Homeost Agents. 2012;26(2 Suppl 1):3S-22S. [PubMed: 23648195].
- Dohan Ehrenfest DM, Andia I, Zumstein MA, Zhang CQ, Pinto NR, Bielecki T. Classification of platelet concentrates (Platelet-Rich Plasma-PRP, Platelet-Rich Fibrin-PRF) for topical and infiltrative use in orthopedic and sports medicine: current consensus, clinical implications and perspectives. *Muscle ligament tendon j.* 2014;4(1):3–9. [PubMed: 24932440]. [PubMed Central: PMC4049647].
- Dohan Ehrenfest DM, Del Corso M, Diss A, Mouhyi J, Charrier JB. Threedimensional architecture and cell composition of a Choukroun's platelet-rich fibrin clot and membrane. *J Periodontol*. 2010;81(4):546– 55. doi: 10.1902/jop.2009.090531. [PubMed: 20373539].

- Badiee P, Alborzi A, Vojdani R, Shakiba E, Rasouli M, Ravanfar P, et al. Early diagnosis of systemic candidiasis in bone marrow transplant recipients. *Exp Clin Transplant*. 2010;8(2):98–103. [PubMed: 20565365].
- Badiee P, Amirghofran AA, Ghazi Nour M, Shafa M, Nemati MH. Incidence and outcome of documented fungal endocarditis. *Int Cardiovasc Res J*. 2014;8(4):152–5. [PubMed: 25614858]. [PubMed Central: PMC4302502].
- Shelburne SA, Sahasrabhojane P, Saldana M, Yao H, Su X, Horstmann N, et al. Streptococcus mitis strains causing severe clinical disease in cancer patients. *Emerg Infect Dis.* 2014;20(5):762–71. doi: 10.3201/eid2005.130953. [PubMed: 24750901]. [PubMed Central: PMC4012796].
- Mager DL, Ximenez-Fyvie LA, Haffajee AD, Socransky SS. Distribution of selected bacterial species on intraoral surfaces. *J Clin Periodontol*. 2003;30(7):644–54. doi: 10.1034/j.1600-051X.2003.00376.x.
- Eren AM, Borisy GG, Huse SM, Mark Welch JL. Oligotyping analysis of the human oral microbiome. *Proc Natl Acad Sci U S A*. 2014;111(28):E2875-84. doi: 10.1073/pnas.1409644111. [PubMed: 24965363]. [PubMed Central: PMC4104879].
- Esteban-Tejeda L, Malpartida F, Esteban-Cubillo A, Pecharroman C, Moya JS. The antibacterial and antifungal activity of a soda-lime glass containing silver nanoparticles. *Nanotechnol.* 2009;**20**(8):85103. doi: 10.1088/0957-4484/20/8/085103. [PubMed: 19417439].
- Jain J, Arora S, Rajwade JM, Omray P, Khandelwal S, Paknikar KM. Silver nanoparticles in therapeutics: development of an antimicrobial gel formulation for topical use. *Mol Pharm.* 2009;6(5):1388–401. doi: 10.1021/mp900056g. [PubMed: 19473014].
- Gajbhiye M, Kesharwani J, Ingle A, Gade A, Rai M. Fungus-mediated synthesis of silver nanoparticles and their activity against pathogenic fungi in combination with fluconazole. *Nanomed.* 2009;5(4):382–6. doi: 10.1016/j.nano.2009.06.005. [PubMed: 19616127].
- Jun EA, Lim KM, Kim K, Bae ON, Noh JY, Chung KH, et al. Silver nanoparticles enhance thrombus formation through increased platelet aggregation and procoagulant activity. *Nanotoxicol*. 2011;5(2):157–67. doi: 10.3109/17435390.2010.506250. [PubMed: 20822370].
- Dohan Ehrenfest DM, Rasmusson L, Albrektsson T. Classification of platelet concentrates: from pure platelet-rich plasma (P-PRP) to leucocyte- and platelet-rich fibrin (L-PRF). *Trends Biotechnol*. 2009;27(3):158–67. doi: 10.1016/j.tibtech.2008.11.009. [PubMed: 19187989].
- Hassan A, Usman J, Kaleem F, Omair M, Khalid A, Iqbal M. Evaluation of different detection methods of biofilm formation in the clinical isolates. *Brazil J Infect Dis.* 2011;15(4):305–11. [PubMed: 21860999].
- Cieslik-Bielecka A, Dohan Ehrenfest DM, Lubkowska A, Bielecki T. Microbicidal properties of Leukocyte- and Platelet-Rich Plasma/Fibrin (L-PRP/L-PRF): new perspectives. J Biol Regul Homeost Agents. 2012;26(2 Suppl 1):43S-52S. [PubMed: 23648198].
- Pinto NR, Ubilla M, Zamora Y, Del Rio V, Dohan Ehrenfest DM, Quirynen M. Leucocyte-and platelet-rich fibrin (L-PRF) as a regenerative medicine strategy for the treatment of refractory leg ulcers: a prospective cohort study. *Platelets*. 2018;**29**(5):468–75. doi: 10.1080/09537104.2017.1327654. [PubMed: 28727481].
- 17. Peck MT, Marnewick J, Stephen LXG, Singh A, Patel N, Majeed A. The

use of leukocyte-and platelet-rich fibrin (L-PRF) to facilitate implant placement in bone-deficient sites: a report of two cases: case report. *South Africa Dent J.* 2012;**67**(2):54–6. [PubMed: 23189893].

- Marenzi G, Riccitiello F, Tia M, di Lauro A, Sammartino G. Influence of Leukocyte- and Platelet-Rich Fibrin (L-PRF) in the Healing of Simple Postextraction Sockets: A Split-Mouth Study. *Biomed Res Int.* 2015;2015;369273. doi: 10.1155/2015/369273. [PubMed: 26273612]. [PubMed Central: PMC4529911].
- Haghgoo R, Ataie M, Rezvani MB, Labibzadeh A. [Evaluation of the effect of conventional varnish and nano-silver containing varnish on microleakage of amalgam restoration]. *Daneshvar Med.* 2013;**20**(104):85–90. Persian.
- Alt V, Bechert T, Steinrucke P, Wagener M, Seidel P, Dingeldein E, et al. An in vitro assessment of the antibacterial properties and cytotoxicity of nanoparticulate silver bone cement. *Biomaterials*. 2004;25(18):4383–91. doi: 10.1016/j.biomaterials.2003.10.078. [PubMed: 15046929].
- Zhang XF, Liu ZG, Shen W, Gurunathan S. Silver nanoparticles: synthesis, characterization, properties, applications, and therapeutic approaches. *Int J Mol Sci.* 2016;17(9). doi: 10.3390/ijms17091534. [PubMed: 27649147]. [PubMed Central: PMC5037809].
- Kim JS, Kuk E, Yu KN, Kim JH, Park SJ, Lee HJ, et al. Antimicrobial effects of silver nanoparticles. *Nanomed*. 2007;3(1):95–101. doi: 10.1016/j.nano.2006.12.001. [PubMed: 17379174].
- Badiee P, Zareifar S, Haddadi P, Jafarian H. Incidence of fungal infections in pediatric patients with hematologic neoplasms. *Archiv Pediatr Infect Dis.* 2017;5(3). e41317. doi: 10.5812/pedinfect.41317.
- Haddadi P, Zareifar S, Badiee P, Alborzi A, Mokhtari M, Zomorodian K, et al. Yeast colonization and drug susceptibility pattern in the pediatric patients with neutropenia. *Jundishapur J Microbiol*. 2014;7(9). e11858. doi: 10.5812/jjm.11858. [PubMed: 25485060]. [PubMed Central: PMC4255375].
- Singh A, Verma R, Murari A, Agrawal A. Oral candidiasis: An overview. J Oral Maxillofac Pathol. 2014;18(Suppl 1):S81-5. doi: 10.4103/0973-029X.141325. [PubMed: 25364186]. [PubMed Central: PMC4211245].
- 26. Dehghani Nazhvani A, Haddadi P, Badiee P, Malekhoseini SA, Jafarian H. Antifungal effects of common mouthwashes on candida strains colonized in the oral cavities of liver transplant recipients in south Iran in 2014. *Hepat Mon*. 2016;16(1). doi: 10.5812/hepatmon.31245.
- Badiee P, Alborzi A. Assessment of a real-time PCR method to detect human non-cryptococcal fungal meningitis. *Arch Iran Med.* 2011;14(6):381-4. [PubMed: 22039841].
- Badiee P, Alborzi A, Shakiba E, Farshad S, Japoni A. Susceptibility of Candida species isolated from immunocompromised patients to antifungal agents. *East Mediterr Health J.* 2011;17(5):425–30. [PubMed: 21796956].
- Lu HZ, Weng XH, Zhu B, Li H, Yin YK, Zhang YX, et al. Major outbreak of toxic shock-like syndrome caused by Streptococcus mitis. *J Clin Microbiol*. 2003;41(7):3051–5. [PubMed: 12843042]. [PubMed Central: PMC165286].
- Papaioannou W, Gizani S, Haffajee AD, Quirynen M, Mamai-Homata E, Papagiannoulis L. The microbiota on different oral surfaces in healthy children. Oral Microbiol Immunol. 2009;24(3):183–9. doi: 10.1111/j.1399-302X.2008.00493.x.