Evaluation of the Chemical Compounds and Antioxidant and Antimicrobial Activities of the Leaves of *Eichhornia Crassipes* (Water Hyacinth)

Rudابه Rufchaei¹, Mahdi Abbas-Mohammadi^{2,*}, Alireza Mirzajani¹ and Shiva Nedaei³

¹Iranian Fisheries Science Research Institute, Inland Waters Aquaculture Research Center, Agricultural Research Education and Extension Organization (AREEO), Bandar Anzali, Iran

²Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, Iran

³Department of Fisheries Science, Faculty of Marine Science and Technology, University of Hormozgan, Bandar Abbas, Iran

*Corresponding author: Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, Iran. Email: mam814@uowmail.edu.au

Received 2020 March 28; Revised 2021 May 15; Accepted 2021 May 17.

Abstract

Background: Water hyacinth (WH) is an aquatic weed and one of the most productive plants on earth, causing serious environmental problems. Herein, some nutritional and phytochemical constituents of WH were investigated.

Methods: Chemical analysis of *Eichhornia Crassipes* was carried out to determine total ash, humidity, crude protein, fat, fiber, and carbohydrate contents. Total phenolic and total flavonoid contents of the hydro-methanolic and aqueous extracts of the plant were determined using the Folin Ciocalteu and aluminum chloride colorimetric methods, and HPLC was performed to quantify eight phenolic compounds. The antioxidant and antimicrobial activities of the extracts were also evaluated.

Results: The dry matter, total ash, crude protein, crude fiber, nitrogen-free extract, and ether extract contents of WH constituted 9.4, 12.9, 24, 11.5, 49.9, and 1.7%, respectively. The total phenolic contents of the hydro-methanolic and aqueous extracts were 491.2 ± 31.9 and 258.3 ± 10.8 mg gallic acid equivalents/g of dried extract, respectively. The total flavonoid content of the hydro-methanolic extract (76.8 ± 7.8) was higher than that of the aqueous extract (46.1 ± 6). Ferulic acid was found to be the most abundant phenolic compound in both extracts. The antioxidant activities of the hydro-methanolic and aqueous extracts were determined to be 221.52 and 97.07 mg ascorbic acid equivalent/g dry weight, respectively. The aqueous and hydro-methanolic extracts showed the highest antibacterial activity against *Escherichia coli*, *Pseudomonas aeruginosa*, respectively.

Conclusions: In conclusion, the present study indicated the applicability of WH as a natural source of antioxidants and antimicrobial agents.

Keywords: Nutritive Value, *Eichhornia Crassipes*, Phenolic Acids, Antioxidant, Antibacterial

1. Background

Water hyacinth (*Eichhornia Crassipes*), belonging to the family of Pontederiaceae, is a wild freshwater fern. This plant is native to South America but has been naturalized in many tropical and subtropical regions of the world. Water hyacinth is considered the world's most rapidly growing and spreading aquatic plant with congested growth, which is also a source of biomass. It is a species with a great ornamental value, used in gardening because of the beauty of its foliage and flowers, but at the same time, water hyacinth is the most important aquatic weed of the world as well (1).

At the first glance, when researchers from Guilan University, Iran, observed water hyacinth in their research tour in Aynak lagoon, it was considered a threat because of

its aggressive expanding on water, leading to the disappearing of native vegetation, decreasing of water transfer capacity, and also fishing restrictions. The plant was even introduced as a toxic bioterrorism agent. However, the plant's beauty of flowers led to its fast spread across Guilan province as a decorative plant (2). Although water hyacinth is often seen as a weed responsible for serious problems in navigation, irrigation, and power generation, several unique properties advocate useful applications for the plant. For example, the plant is efficient in removing toxic metals from water (3). If the chemical composition and nutritive value of *E. crassipes* can be explored, farmers can utilize this plant as an unconventional feed for their livestock to minimize feed costs and maximize production (4).

Water hyacinth is a source of chemicals with medicinal functions (5-7). The leaf extract of this plant contains flavonoids, alkaloids, tannins, and phenols, which have biological activities such as antiviral, antifungal, antitumor, and antibacterial effects (8-10). In addition, water hyacinth is rich in oxidative enzymes and non-enzymatic antioxidant systems (6), as well as wound healing effects and anticancer activities (8). The antioxidative properties of *E. crassipes* make this plant an alternative and convenient source of antioxidants (11).

Medicinal plants are rich resources of different bioactive compounds that can be used to treat several diseases by modulating various enzymes (12, 13). Plant-derived products have been used as disease controlling agents and reported to have less toxicity and fewer unwanted effects on the environment. An alternative approach for identifying new and innovative medicinal products is to search for such agents with antimicrobial and antifungal activities in medicinal plants, as well as in common weeds, which are also sources for several antimicrobial and antifungal compounds (14, 15).

2. Objectives

Many studies have already investigated the chemical composition of water hyacinth growing all over the world. There is evidence that the biosynthesis patterns of secondary metabolites in different populations of a single plant species can vary in different geographical regions and in response to environmental changes (16). Despite this, there is no information regarding the phytochemical structure of the water hyacinth species growing in West Asia. Therefore, in pursuing our research on water hyacinth (17, 18), we aimed to discover the chemical composition and nutrients of the water hyacinth growing in the Guilan province of Iran. We also determined the antioxidant and antimicrobial activities of the hydro-methanolic and aqueous extracts of the plant.

3. Methods

3.1. Chemicals

Methanol and ethanol (technical grade, purified by distillation) were used for the extraction procedure and total flavonoid determination, respectively. Acetonitrile (high-performance liquid chromatography (HPLC) grade), trifluoroacetic acid, and the Folin Ciocalteu reagent were purchased from Merck Co. (Germany). Phenolic compounds (gallic acid, catechin, chlorogenic acid, vanillic acid, P-coumaric acid, ferulic acid, rutin, and quercetin, as

well as the Fe^{3+} -2,4,6-tri (2-pyridyl)-s-triazine (TPTZ) complex, $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, aluminum chloride, potassium acetate, and sodium carbonate were purchased from the Sigma Chemical Co. (St. Louis, MO, USA). The standard strains of *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Aeromonas hydrophila*, and *Streptococcus iniae* were also provided by the Iranian Research Organization for Science and Technology and the Microbiology Research Laboratory of Guilan University (Table 1).

Table 1. The Specifications of the Organisms Studied

Microbial species	Source	Gram
<i>Staphylococcus aureus</i>	6538. Iranian Research Organization for Science and Technology	Positive cocci
<i>Escherichia coli</i>	8739. Iranian Research Organization for Science and Technology	Negative coccobacilli
<i>Pseudomonas aeruginosa</i>	9027. Iranian Research Organization for Science and Technology	Negative coccobacilli
<i>Aeromonas hydrophila</i>	7966. Guilan University microbiology lab	Negative bacilli
<i>Streptococcus iniae</i>	29177. Guilan University microbiology lab	Positive cocci

3.2. Plant Materials

The leaves of *E. crassipes* were collected from Anzali lagoon, in the Caspian Sea near Bandar-e-Anzali, the northern Iranian province of Guilan, in Oct 2016 and identified in the Faculty of Natural Resources of Guilan University.

3.3. Extraction and Isolation

The leaves of the plant were dried at room temperature and ground into a fine powder with the particle mesh size of 1-mm using a mechanical grinder. Two parts (10 g each) of the powdered plant were extracted at room temperature using the maceration procedure using two solvents: purified water (3×100 mL) and analytical grade methanol 80% (3×100 mL). The extracts of each solvent were combined and allowed to evaporate and dry under reducing pressure at 45°C to obtain less solventaqueous and hydro-methanolic extracts. The yield of extraction was calculated as 1.2 and 2.9% for the aqueous and hydro-methanolic extracts, respectively.

3.4. Chemical Analyses

Chemical analyses of *E. crassipes* were carried out in triplicate to determine total ash, humidity, crude protein, crude fat, crude fiber, and crude carbohydrate contents. The total ash content of the plant was measured utilizing an electrical furnace (Heraeus K1252 muffle furnace) at

550°C for five hours. A Binder oven (WTB Binder- Drying oven in Barneveld, the Netherlands) was used to measure the humidity of the plant at 150°C for 12 hours. Protein content was determined using Kjeltac TM 2300 Analyzer (FOSS, Hilleroed, Denmark). Crude fat was determined using a soxtec 1043 system. Crude fiber and carbohydrate contents were also estimated using an enzymatic-gravimetric AOAC method by measuring the loss in ignition of dried lipid-free residues after digestion with 1.25% H₂SO₄ and 1.25% NaOH in a Fiber cap 2021/2023 system (Foss Tecator, Sweden) (19).

3.5. Determination of Total Phenolic Content

The concentrations of phenolic compounds in the plant extracts were determined using the spectrophotometric Folin Ciocalteu method (20). Crude extracts were prepared at the concentration of 10 mg. mL⁻¹ in water. The reaction mixture was prepared by mixing 0.1 mL of the extracts, 0.1 mL of 50% the Folin Ciocalteu reagent dissolved in water, 2 mL of 2% NaHCO₃, and 2.8 mL of distilled water. Then, the samples were incubated at 25°C for 30 min, and absorbance was measured at 750 nm. Each analysis was performed in triplicate. Gallic acid was used to draw a standard curve, and based on the sample's absorbance, the concentration of phenolic compounds was expressed as gallic acid equivalent (mg of GA/g of extract).

3.6. Determination of Total Flavonoid Content

Total flavonoid content was determined by the aluminum chloride colorimetric method, with slight modifications (21). For this purpose, aliquots were prepared by dissolving 10 mg of the extracts in 5 mL deionized water. Then 0.5 mL of the prepared extracts was mixed with 1.5 mL of EtOH 95%, 100 µL of AlCl₃ (10%), 100 µL of 1M potassium acetate, and 2.8 mL deionized water, followed by incubation at room temperature for 40 min. After that, absorbance was read at 415 nm. Quercetin was used as a standard. Total flavonoid content was expressed as mg quercetin equivalents/g dry mass. The experiment was performed in triplicate for both extracts.

3.7. Quantification of Phenolic Compounds by HPLC Analysis

Eight phenolic compounds, including gallic acid, catechin, chlorogenic acid, vanillic acid, P- coumaric acid, ferulic acid, rutin, and quercetin, were quantified using a Knauer HPLC system and a SunFire C18 column (250 mm × 4.6 mm, 5 µm), applying the Knauer photodiode array detector (PDA; K-2800 model). Injection was performed through a 3900 Smartline autosampler injector equipped with a loop (100 µL). The quantification of phenolic compounds was performed based on an optimized

HPLC method described by Alirezalu et al., (22) with some modifications. The effects of different parameters, including the mobile phase, column, flow rate, and solvent ratio, were assessed to optimize chromatographic conditions. According to the results obtained, the column was eluted at a flow rate of 0.5 mL/min with water + 0.02% trifluoroacetic acid (solvent A) and acetonitrile + 0.02% trifluoroacetic acid (solvent B); the following gradient was set as: 0 - 20 min, from 20% B to 40% B; 20 - 40 min, 40% B; 40 - 42 min, from 40% B to 100% B; 42 - 48 min, 100% B; 48 - 50 min, from 100% B to 20% B (i.e., return to the initial condition), and 50 - 55 min; 20% B.. The temperature of the column was kept at 25°C. Chromatograms were recorded at four different wavelengths, including 254, 273, 310, and 326 nm (according to the λ_{max} of the evaluated compounds), with a run time of 55 min. Data acquisition and integration were performed by EZChrome Elite software. The quantification of phenolic compounds was performed based on calibration via standards and peak-area measurements. For drawing calibration curves, the standard of each phenolic compound was weighed accurately (1 mg) and dissolved in 1 mL of MeOH (HPLC grade) to prepare a stock solution. The stock solutions were then diluted by MeOH to obtain a series of concentrations. Next, 20 µL of each solution was injected into the HPLC device. Based on the standard solutions, mean areas (n = 3) were determined and plotted against the concentration to develop calibration equations (23, 24).

3.8. Evaluation of Antioxidant Activity

The antioxidant capacity of the extracts was spectrophotometrically estimated following the procedure of Benzie and Strain (25). This method is based on reducing the colorless Fe³⁺-2,4,6-tri (2-pyridyl)-s-triazine (TPTZ) complex to the blue colored Fe²⁺-tripyridyltriazine complex by the action of electron donating antioxidants at low pH. The FRAP reagent was prepared as a solution containing 25 mL of 300 mM acetate buffer (pH = 3.6), 2.5 mL of 10 mM TPTZ solution in 40 mM HCl, and 2.5 mL of 20 mM FeCl₃.6H₂O. To determine antioxidant capacity, 100 µL of the extract solutions was mixed with 3.0 mL of freshly prepared FRAP reagent and incubated at 37°C for 10 min. Absorbance was recorded at 593 nm against a blank sample containing 100 µL of the solvent. The FRAP value was calculated from the calibration curve prepared according to FeSO₄.7H₂O standard solutions and expressed as mmol Fe²⁺/g dry weight.

3.9. Antimicrobial Assays

The antimicrobial activity of the extracts was evaluated by measuring bacterial growth inhibition zones by

the well-diffusion method, as described by the Clinical Laboratory Standard Institute (CLSI) (26), against five standard strains, including *S. aureus*, *E. coli*, *P. aeruginosa*, *A. hydrophila*, and *S. iniae*. Briefly, 0.1 mL of the diluted inoculum (105 CFU/mL) of the test strains was swabbed on nutrient agar in plates. Wells with a 6-mm diameter were punched into the agar surface via a sterile cork borer. According to our previous study (27), 50 μ L of the plant extracts in a serial dilution of 5, 20, 100, and 400 mg.mL⁻¹ was added to the wells created in the plates. The plates were incubated at 37°C for 24 hr. Antibacterial activity was determined by measuring the diameter of the zone of inhibition (ZOI) surrounding wells. Chloramphenicol standard antibiotic disc was used as the positive control, and the wells containing 20 μ L DMSO were used as the negative control.

The broth microdilution method was also carried out to determine the minimum inhibitory concentration (MIC) of the extracts according to the standard protocol recommended by Eloff (28). In brief, a serial dilution of each extract was prepared in a concentration range of 32 to 1024 mg. mL⁻¹ in sterile 96 wells containing Mueller-Hinton broth medium, using DMSO as the solvent of the extracts at a final concentration of lower than 1% (v/v). Suspensions from the test strains were prepared from freshly cultured microorganisms in sterile normal saline, adjusted to the turbidity of 0.5 McFarland standard. The suspensions were further diluted (1:100) by the mentioned sterile broth medium and then added to the trays containing a serial dilution of the extracts and 0.5×10^6 CFU/mL bacteria per well. The trays containing serial dilutions were incubated at 37°C for 24 h. The incubation of aquatic strains was performed at 28°C. Finally, MIC values were recorded as the lowest concentrations that could inhibit the visible growth of microorganisms. The resazurine reagent was used for the extracts forming turbidity after being dissolved to characterize growth from no-growth wells. For determining the minimum bactericidal concentration (MBC), 100 μ L of each no-growth well identified in MIC determination experiments was cultured in Mueller Hinton Broth, and MBC values, as the lowest concentration killing 99.9% of the microorganism tested, were recorded after 24 h of incubation at 37°C. Each experiment was conducted in triplicate.

3.10. Statistical Analysis

All experiments were carried out in triplicate. The results obtained were expressed as average $\pm$ standard error of mean. The Shapiro-Wilk test was used to evaluate the normality of the data. The data were analyzed in SPSS 16.0 software by the independent *t*-test. Pearson correlation coefficient was utilized to investigate the correlation of an-

tioxidant activity with the total phenol and flavonoid contents of the plant.

4. Results

In the present study, the nutritional and phytochemical constituents of water hyacinth were investigated. The values of dry mass, total ash, nitrogen-free extract, crude protein, ether extract, and crude fiber were obtained 9.4, 12.9, 49.9, 24, 1.7, and 11.5%, respectively (Table 2).

Table 2. The Chemical Analysis of Water Hyacinths

Parameters	Percent (Based on DM)
Dry matter	9.4
Crude protein	24
Ether extract	1.7
Crude fiber	11.5
Ash	12.9
Nitrogen free extract	49.9

The aqueous and hydro-methanolic extracts of *E. crassipes* were used to investigate the phytochemical constituents of the plant. The yields of extraction were 12 and 29% for the hydro-methanolic and aqueous extracts, respectively. The total phenol contents of the hydro-methanolic and aqueous extracts, expressed as gallic acid equivalent, were 491.2 ± 31.9 and 258.3 ± 10.8 mg gallic acid equivalents/g of dried extract, respectively. The total flavonoid contents of the hydro-methanolic and aqueous extracts expressed as mg quercetin equivalents/g of dried extracts, were 76.8 ± 7.8 and 46.1 ± 6 , respectively, showing a significant difference.

The amounts of seven phenolic compounds, including gallic acid, catechin, chlorogenic acid, vanillic acid, p-coumaric acid, ferulic acid, rutin, and quercetin, were simultaneously analyzed by HPLC. The UV patterns of standard analyses were compared with those related to the two extracts of water hyacinth. In addition, spiking of each standard phenolic to the extracts was applied to exactly assign the signal of phenolics in the HPLC chromatograms. The HPLC chromatograms of standard phenolic compounds have been shown in Figure 1, indicating an appropriate resolution for different signals. In addition, the HPLC chromatograms of the aqueous (A) and hydro-methanolic (B) extracts have been indicated in Figure 2. In this regard, three compounds (rutin, quercetin, and vanillic acid) were quantified at 254 nm; gallic acid and catechin were measured at 273 nm; p-coumaric acid was analyzed at 310 nm, and finally, ferulic acid and chlorogenic acid were assessed at 326 nm. Figure 3 reveals the overlaid

UV-visible spectra of the evaluated compounds in the extracts and standard samples. A good agreement was observed between the spectra of the standards and those related to the extracts' constituents, confirming the peak purity and accurate identification of each signal. Table 3 summarizes the contents of phenolic compounds in the hydro-methanolic and aqueous extracts analyzed in this study. Ferulic acid was found to be the most abundant phenolic compound in both the hydro-methanolic and aqueous extracts (2.28 and 2.54 mg/g of dried matter, respectively), showing a slightly higher value in the aqueous extract. Other phenolic contents had significantly variable quantities in the two extracts (Table 3). Chlorogenic acid and quercetin were two other major phenolic compounds of the aqueous extract (0.81 and 0.55 mg/g of dried matter, respectively). In the hydro-methanolic extract, quercetin, catechin, and gallic acid were found as other main phenolic compounds with the values of 1.31, 0.7, and 0.62 mg/g, respectively.

Table 3. Phenolic Compounds in Two Extracts of Water Hyacinth ^{a, b}

Phenolic Compounds	Aqueous Extract	Hydro-Methanolic Extract
	mg/g of Dried Mass	
Gallic acid	0.260 ± 0.010 ^A	0.620 ± 0.010 ^B
Catechin	0.270 ± 0.005 ^A	0.700 ± 0.004 ^B
Chlorogenic acid	0.810 ± 0.005 ^B	0.230 ± 0.030 ^A
Vanillic acid	0.120 ± 0.005 ^A	0.210 ± 0.004 ^B
p- coumaric acid	0.120 ± 0.020 ^A	0.100 ± 0.009 ^A
Ferulic acid	2.540 ± 0.020 ^B	2.280 ± 0.010 ^A
Rutin	0.150 ± 0.020 ^A	0.320 ± 0.010 ^B
Quercetin	0.550 ± 0.000 ^A	1.310 ± 0.010 ^B

^a Values are expressed as mean ± SE.

^b Data in the same row with different capital letters (A - B) show significant differences ($P < 0.05$).

In the present research, FRAP, as a commonly accepted assay, was used to evaluate the antioxidant activity of the extracts. This assay was selected considering the fact that the antioxidant activity of plant derivatives depends on their reducing capacity (29). The FRAP values were obtained as 221.52 and 97.07 mg ascorbic acid equivalent/g of dry weight of the hydro-methanolic and aqueous extracts, respectively.

Pearson correlation coefficients were used to investigate the possible correlation between the antioxidant activity and phytochemical constituents of the extracts analyzed (Table 4). In both extracts, antioxidant activity significantly correlated with the total phenolic ($P = 0.01$) and total flavonoid ($P < 0.05$) contents.

The antibacterial activity of the aqueous and hydro-methanolic extracts was investigated using the agar well-diffusion assay against five standard strains, including *E. coli*, *S. aureus*, and *P. aeruginosa*, as well as two aquatic pathogens (i.e., *A. hydrophila*, and *S. iniae*) for the first time. Table 5 indicates the diameter of the zone of inhibition for the extracts at different concentrations against the five bacterial strains. According to the results obtained, the aqueous extract was the most active against *E. coli* with an inhibition zone of 15.3 mm at the concentration of 400 mg/mL. At the same concentration, the antibacterial activity of the methanolic extract was the highest against *S. iniae*, with an inhibition zone of 12.3 mm.

The aqueous extract showed the highest antibacterial activity, with the MIC and MBC values of 64 and 128 mg.mL⁻¹, against *E. coli*. Furthermore, the methanolic extract delivered the best antibacterial activity against *S. iniae* with the MIC and MBC values of 64 and 128 mg/mL, respectively. The most significant antibacterial resistance was observed for *P. aeruginosa* in exposure to both extracts with the MIC and MBC values of 256 and 512 mg/mL, respectively. Regarding the two other strains tested (i.e., *A. hydrophila* and *S. aureus*), both the extracts showed moderate antibacterial activity as indicated by respective MIC and MBC values (Table 6).

5. Discussion

The plant species studied (water hyacinth) had a high value of crude protein compared to other plants; however, its crude fiber content was lower compared to previously reported values. These variations can be due to different environmental and growing conditions. The contents of other constituents were in well agreement with previously reported values (30-32).

The total phenol and total flavonoid contents of plant extracts partly depend on the polarity of the solvent used in extraction. The high solubility of phenols and flavonoids in polar solvents leads to a high extraction yield for these compounds (33). In the present experiment, the aerial parts of the plant were extracted using two polar solvents (distilled water and hydro-methanol). According to the results obtained, the total phenol and flavonoid contents of the hydro-methanolic extract were higher compared with the aqueous extract. Similar findings were reported by Ho et al. and Rorong et al. (9, 34, 35).

Phenolic compounds are well-known for their potent antioxidant properties and are commonly found in fruits, vegetables, and herbs (36, 37). So, the high antioxidant activity of the hydro-methanolic extract in this research can be attributed to its high amounts of polyphenols and flavonoids. Gallic acid and quercetin, as powerful natural antioxidants, were identified in the plant extracts as-

Figure 1. The HPLC chromatograms of standard phenolic compounds

Table 4. The Correlation Between Total Phenol and Total Flavonoid Contents and Antioxidant Activity

Antioxidant Activity	Total Phenols		Total Flavonoids	
	Hydro-Methanolic	Aqueous	Hydro-Methanolic	Aqueous
P value	0.014	0	0.018	0.003
R	0.9 ^a	0.99 ^b	0.889 ^a	0.959 ^b

^a Significant at $P < 0.05$.

^b Significant at $P < 0.01$.

Table 5. The Antimicrobial Activity of *E. crassipes* Hydro-Methanolic and Aqueous Extracts Against Different Bacteria Based on the Well-Diffusion Method^{a, b}

Microorganisms	Inhibition Zone Diameter								Control	
	Aqueous Extract (mg/mL)				Hydro-Methanolic Extract (mg/mL)				Negative ^c	Positive ^d
	5	20	100	400	5	20	100	400	(-)	(+)
<i>E. coli</i>	8.0 ± 0.5 ^E	9.0 ± 0.5 ^D	11.6 ± 0.3 ^C	15.3 ± 1.0 ^B	6.0 ± 0.0	7.0 ± 0.5 ^D	9.6 ± 0.3 ^C	11.6 ± 0.3 ^B	6.0 ± 0.0	22.0 ± 0.0 ^A
<i>S. aureus</i>	6.0 ± 0.0	4.0 ± 0.5 ^D	10.3 ± 0.6 ^C	11.0 ± 1.0 ^B	7.0 ± 0.3 ^E	8.0 ± 0.6 ^D	11.0 ± 0.5 ^C	11.3 ± 0.3 ^B	6.0 ± 0.0	25.0 ± 0.0 ^A
<i>P. aeruginosa</i>	6.0 ± 0.0	6.0 ± 0.0	6.0 ± 0.0	9.6 ± 0.3 ^A	6.0 ± 0.0	6.0 ± 0.0	6.0 ± 0.0	10.3 ± 0.6 ^A	6.0 ± 0.0	11.0 ± 0.0 ^A
<i>A. hydrophila</i>	6.0 ± 0.0	6.0 ± 0.0	8.3 ± 1.0 ^C	11.3 ± 2.0 ^B	6.0 ± 0.0	6.0 ± 0.0	10.0 ± 1.0 ^C	11.6 ± 0.3 ^B	6.0 ± 0.0	30.0 ± 0.0 ^A
<i>S. iniae</i>	6.0 ± 0.0	6.0 ± 0.0	9.3 ± 0.3 ^C	10.6 ± 0.3 ^B	8.0 ± 0.5 ^E	10.0 ± 1.0 ^C	9.0 ± 0.5 ^C	12.3 ± 0.6 ^B	6.0 ± 0.0	25.0 ± 0.0 ^A

^a Values are expressed as mean ± SE.

^b Data in the same row with different capital letters (A - E) show significant differences ($P < 0.05$).

^c Standard antibiotic discs (30 µg/mL).

^d DMSO.

sessed in the present study. The ability of gallic acid to form hybrid compounds with other phenols such as caffeic acid increases its antioxidant properties (38). In addition, quercetin has been shown to have a more potent antioxidant activity than aglycone-binding flavonoids such as rutin due to the lack of an aglycon site in its structure (39). In this study, HPLC analysis showed a high amount of quercetin in the hydroalcoholic extract of water hyacinth (1.31 mg/g dry extract). However, quercetin had a low concentration in the aqueous extract. So, it is possible that

the higher In addition, spiking of each standard phenolic to the extracts was applied to exactly assign the signal of phenolics in the HPLC chromatograms. Power of the hydro-alcoholic extract compared to the aqueous extract can be due to the higher concentration of this flavonoid. Rutin, in synergism with other antioxidants, has shown the ability to eliminate free radicals and augment protection against fat oxidation (40).

Regarding the phenolic compounds of water hyacinth observed in the present study, a similar profile has also

Figure 2. The HPLC chromatograms of both the aqueous (A, right) and hydro-methanolic (B, left) extracts at four evaluated wavelengths.

been reported in other studies (9, 41). However, there is limited information concerning the quantity of phenolic compounds in water hyacinth. In the only research that we found on water hyacinth (42), the content of quercetin, chlorogenic acid, and catechin was estimated to be zero, which is in contrast to our results. These contradictory re-

sults may be due to different extraction methods or the effects of ecological parameters (43). Further research is required to accurately quantify and qualify the secondary metabolites of water hyacinth (5, 41). Polyphenol and flavonoid contents of the aqueous and methanolic extracts of 31 marsh medicinal plants were measured in a study

Figure 3. The overlaid UV-visible spectra of the phenolic compounds quantified in the plant extracts and standard samples.

in Taiwan, indicating the predominance of phenolic compounds in the aqueous extracts of most of the plants. However, in water hyacinth, the amounts of total phenols and flavonoids were higher in the methanolic than in aqueous extract (34).

Many researchers have also evaluated the antimicrobial activities of various plant extracts. In one study, the

methanolic extract of water hyacinth significantly reduced the growth of *Aspergillus niger* (44). A significant antimicrobial activity was reported for the ethanolic, methanolic, and aqueous extracts of the roots and leaves of water hyacinth against different bacterial and fungal strains. Among these, the aqueous extract showed a more potent activity, and also, leaves delivered a higher antimicrobial

Table 6. The MBC and MIC Values of the Hydro-Methanolic and Aqueous Extracts of *E. crassipes* Against the Microorganisms Tested

Extracts/Yield (%) / Microorganism	MIC (mg/mL)	MBC mg/mL)
Hydro-Methanolic		
12		
<i>Escherichia coli</i>	128	256
<i>Staphylococcus aureus</i>	128	256
<i>Pseudomonas aeruginosa</i>	> 256	512
<i>Aeromonas hydrophila</i>	256	> 256
<i>Streptococcus iniae</i>	64	128
Aqueous		
29		
<i>Escherichia coli</i>	64	128
<i>Staphylococcus aureus</i>	128	256
<i>Pseudomonas aeruginosa</i>	> 256	512
<i>Aeromonas hydrophila</i>	256	512
<i>Streptococcus iniae</i>	128	256

activity than roots (45). Likewise, water hyacinth extract was reported to have activity against *S. aureus*, *E. coli*, *Penicillium*, and *A. niger*; however, this activity depended on pH, concentration, and duration of exposure (46). The antimicrobial properties of polyphenols have been widely evaluated and confirmed in previous reports (47, 48). Although the mechanisms of the antibacterial actions of phenolic compounds are not yet fully deciphered, polyphenols have been suggested to exert their antibacterial effects via either direct killing of bacteria, the synergistic activation of antibiotics, and attenuation of the pathogenicity of bacteria (49). More importantly, flavonoids have been shown to inactivate efflux pumps, destabilize cytoplasmic membrane, and inhibit β -lactamases and topoisomerase and, in this way, prevent the development of antibiotic resistance in bacteria (48). Nevertheless, the antibacterial effects of polyphenols, in addition to the type of polyphenol, can also be influenced by other factors. For instance, the same polyphenol may be effective against one Gram-positive (or Gram-negative) strain and ineffective against the other, indicating strain-dependent effects.

5.1. Conclusion

The present study revealed high amounts of biologically active phenols and flavonoids in the hydro-methanolic extract of *E. crassipes*. The high content of polyphenols in this plant extract could justify its potent antioxidant activity. Our results suggest that *E. crassipes* can be beneficial as an antimicrobial agent, especially against Gram-negative bacteria, and an available natural source for antioxidants and antimicrobial agents.

Footnotes

Authors' Contribution: R Rufchaei, study supervision; M Abbas-Mohammadi acquisition of data + analysis and interpretation of data + drafting the manuscript; A Mirzajani, study concept and design; S Nedaie, acquisition of data + statistical analysis.

Conflict of Interests: Authors declare that there is no conflict of interest.

Funding/Support: This research was financially supported by the Guilan provincial government and documented by the Agriculture, Research, Education, and Extension Organization.

References

1. Lancar L, Krake K. *Aquatic weeds & their management*. New Delhi, India: International Commission on Irrigation and Drainage; 2002.
2. Mozaffarian V, Yaghoubi B. New record of Eichhornia crassipes (Water Hyacinth) from north of Iran. *Rostaniha*. 2016;**16**(2):208–11.
3. Dandelot S, Robles C, Pech N, Cazaubon A, Verlaque R. Allelopathic potential of two invasive alien Ludwigia spp. *Aquat Bot*. 2008;**88**(4):311–6. doi: [10.1016/j.aquabot.2007.12.004](https://doi.org/10.1016/j.aquabot.2007.12.004).
4. Malik A. Environmental challenge vis a vis opportunity: The case of water hyacinth. *Environ Int*. 2007;**33**(1):122–38. doi: [10.1016/j.envint.2006.08.004](https://doi.org/10.1016/j.envint.2006.08.004). [PubMed: [17010439](https://pubmed.ncbi.nlm.nih.gov/17010439/)].
5. Jayanthi P, Lalitha P, Shubashini KS. Phytochemical investigation of the extracts of Eichhornia crassipes and its solvent fractionates. *J Pharm Res*. 2011;**4**(5):1405–6.
6. Lalitha TP, Jayanthi P. Preliminary studies on phytochemicals and antimicrobial activity of solvent extracts of Eichhornia crassipes (Mart.) Solms. *Asian J Plant Sci Res*. 2012;**2**(2):115–22.
7. Joshi M, Kaur S. In vitro evaluation of antimicrobial activity and phytochemical analysis of Calotropis procera, Eichhornia crassipes and Datura innoxia leaves. *Asian J Pharm Clin Res*. 2013;**6**(5):25–8.

8. Ali H, Patel M, Ganesh N, Ahi J. The world's worst aquatic plant as a safe cancer medicine-Antitumor activity on melanoma induced mouse by Eichhornia crassipes: In vivo studies. *J Pharm Res.* 2009;**2**:1365-6.
9. Shanab SM, Shalaby EA, Lightfoot DA, El-Shemy HA. Allelopathic effects of water hyacinth [Eichhornia crassipes]. *PLoS One.* 2010;**5**(10): e13200. doi: [10.1371/journal.pone.0013200](https://doi.org/10.1371/journal.pone.0013200). [PubMed: [20949048](https://pubmed.ncbi.nlm.nih.gov/20949048/)]. [PubMed Central: [PMC2951916](https://pubmed.ncbi.nlm.nih.gov/PMC2951916/)].
10. Kurup AR, Rajan D, Blessen J, Chandran S, Thampatty TAR, Veena VPV. Detailed analysis on phytochemicals, antioxidants, antimicrobial activity of Eichhornia crassipes. *Int J Sci Res.* 2012;**2**(2):17-9. doi: [10.15373/22778179/feb2013/7](https://doi.org/10.15373/22778179/feb2013/7).
11. Bodo R, Azzouz A, Hausler R. Antioxidative activity of water hyacinth components. *Plant Sci.* 2004;**166**(4):893-9. doi: [10.1016/j.plantsci.2003.12.001](https://doi.org/10.1016/j.plantsci.2003.12.001).
12. Abbas-Mohammadi M, Moridi Farimani M, Salehi P, Nejad Ebrahimi S, Sonboli A, Kelso C, et al. Acetylcholinesterase-inhibitory activity of Iranian plants: Combined HPLC/bioassay-guided fractionation, molecular networking and docking strategies for the dereplication of active compounds. *J Pharm Biomed Anal.* 2018;**158**:471-9. doi: [10.1016/j.jpba.2018.06.026](https://doi.org/10.1016/j.jpba.2018.06.026). [PubMed: [29960238](https://pubmed.ncbi.nlm.nih.gov/29960238/)].
13. Subramanian R, Asmawi MZ, Sadikun A. In vitro alpha-glucosidase and alpha-amylase enzyme inhibitory effects of Andrographis paniculata extract and andrographolide. *Acta Biochim Pol.* 2008;**55**(2):391-8. [PubMed: [18511986](https://pubmed.ncbi.nlm.nih.gov/18511986/)].
14. Cushnie TP, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents.* 2005;**26**(5):343-56. doi: [10.1016/j.ijantimicag.2005.09.002](https://doi.org/10.1016/j.ijantimicag.2005.09.002). [PubMed: [16323269](https://pubmed.ncbi.nlm.nih.gov/16323269/)]. [PubMed Central: [PMC7127073](https://pubmed.ncbi.nlm.nih.gov/PMC7127073/)].
15. Karimi Alavijeh P, Karimi Alavijeh P, Sharma D. A study of antimicrobial activity of few medicinal herbs. *Asian J Plant Sci Res.* 2012;**2**(4):496-502.
16. Wen B, Ren S, Zhang Y, Duan Y, Shen J, Zhu X, et al. Effects of geographic locations and topographical factors on secondary metabolites distribution in green tea at a regional scale. *Food Control.* 2020;**110**:106979. doi: [10.1016/j.foodcont.2019.106979](https://doi.org/10.1016/j.foodcont.2019.106979).
17. Rufchaei R, Mirvaghefi A, Hoseinifar SH, Valipour A, Nedaei S. Effects of dietary administration of water hyacinth (Eichhornia crassipes) leaves extracts on innate immune parameters, antioxidant defence and disease resistance in rainbow trout (Oncorhynchus mykiss). *Aquaculture.* 2020;**515**:734533. doi: [10.1016/j.aquaculture.2019.734533](https://doi.org/10.1016/j.aquaculture.2019.734533).
18. Rufchaei R, Nedaei S, Hoseinifar SH, Hassanpour S, Golsan M, Sayad Bourani M. Improved growth performance, serum and mucosal immunity, haematology and antioxidant capacity in pikeperch (Sander lucioperca) using dietary water hyacinth (Eichhornia crassipes) leaf powder. *Aquac Res.* 2020;**52**(5):2194-204. doi: [10.1111/are.15072](https://doi.org/10.1111/are.15072).
19. Indrayan AK, Agrawal P, Rath AK, Shatru A, Agrawal NK, Tyagi DK. Nutritive value of some indigenous plant rhizomes resembling Ginger. *Nat Prod Radiance.* 2009;**8**(5):507-13.
20. Chaouche T, Haddouchi F, Ksouri R, Medini F, El-Haci IA, Boucherit Z, et al. Antioxidant activity profiling by spectrophotometric methods of phenolic extract of Prasium majus L. *Free Radic Antioxid.* 2013;**3**(1):43-6. doi: [10.1016/j.fra.2013.03.004](https://doi.org/10.1016/j.fra.2013.03.004).
21. Meda A, Lamien CE, Romito M, Millogo J, Nacoulma OG. Determination of the total phenolic, flavonoid and proline contents in Burkina Faso honey, as well as their radical scavenging activity. *Food Chem.* 2005;**91**(3):571-7. doi: [10.1016/j.foodchem.2004.10.006](https://doi.org/10.1016/j.foodchem.2004.10.006).
22. Alirezalu A, Ahmadi N, Salehi P, Sonboli A, Alirezalu K, Mousavi Khaneghah A, et al. Physicochemical characterization, antioxidant activity, and phenolic compounds of hawthorn (Crataegus spp.) fruits species for potential use in food applications. *Foods.* 2020;**9**(4). doi: [10.3390/foods9040436](https://doi.org/10.3390/foods9040436). [PubMed: [32260449](https://pubmed.ncbi.nlm.nih.gov/32260449/)]. [PubMed Central: [PMC7230283](https://pubmed.ncbi.nlm.nih.gov/PMC7230283/)].
23. Arabshahi D. *Studies on selected plant extracts with reference to their nutritional and pharmacological characteristics [dissertation]*. Mysore, India: University of Mysore; 2006.
24. Chen XQ, Xiao JB. RP-HPLC-DAD determination of flavonoids: Separation of quercetin, luteolin and apigenin in Marchantia convoluta. *Iran J Pharm Res.* 2010;**4**(3):175-81.
25. Benzie IF, Strain JJ. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol.* 1999;**299**:15-27. doi: [10.1016/S0076-6879\(99\)99005-5](https://doi.org/10.1016/S0076-6879(99)99005-5). [PubMed: [9916193](https://pubmed.ncbi.nlm.nih.gov/9916193/)].
26. Espinel-Ingroff AV, Pfaller MA. Susceptibility test methods: Yeasts and filamentous fungi. *Manual of Clinical Microbiology.* 2006;**2**(9):1972-86.
27. Rufchaei R, Mirzajani A, Zamani H. Evaluation of phenolic compounds of methanolic and aqueous extracts of water hyacinth leaves And its antifungal effect on Candida Albicans. *Third International Conference on Agriculture, Environment, Urban and Rural Development.* Tehran, Iran. CIVILCA; 2019.
28. Eloff JN. A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Med.* 1998;**64**(8):711-3. doi: [10.1055/s-2006-957563](https://doi.org/10.1055/s-2006-957563). [PubMed: [9933989](https://pubmed.ncbi.nlm.nih.gov/9933989/)].
29. Bruno M, Ilardi V, Lupidi G, Quassinti L, Bramucci M, Fiorini D, et al. Composition and biological activities of the essential oil from a Sicilian accession of Prangos ferulacea (L.) Lindl. *Nat Prod Res.* 2021;**35**(5):733-43. doi: [10.1080/14786419.2019.1598996](https://doi.org/10.1080/14786419.2019.1598996). [PubMed: [30957559](https://pubmed.ncbi.nlm.nih.gov/30957559/)].
30. Abdelhamid AM, Gabr AA. Evaluation of water hyacinth as a feed for ruminants. *Archiv für Tierernährung.* 2009;**41**(7-8):745-56. doi: [10.1080/17450399109428519](https://doi.org/10.1080/17450399109428519).
31. El-Sayed AFM. Effects of fermentation methods on the nutritive value of water hyacinth for Nile tilapia Oreochromis niloticus (L.) fingerlings. *Aquaculture.* 2003;**218**(1-4):471-8. doi: [10.1016/S0044-8486\(02\)00252-1](https://doi.org/10.1016/S0044-8486(02)00252-1).
32. Hossain E, Sikder H, Kabir H, Sarma SM. Nutritive value of water hyacinth (Eichhornia crassipes). *Online J Anim Feed Res.* 2015;**5**(2):40-4.
33. Mohsen SM, Ammar ASM. Total phenolic contents and antioxidant activity of corn tassel extracts. *Food Chem.* 2009;**112**(3):595-8. doi: [10.1016/j.foodchem.2008.06.014](https://doi.org/10.1016/j.foodchem.2008.06.014).
34. Ho YL, Huang SS, Deng JS, Lin YH, Chang YS, Huang GJ. In vitro antioxidant properties and total phenolic contents of wetland medicinal plants in Taiwan. *Bot Stud.* 2012;**53**(1).
35. Rorong JA, Sudiarso S, Prasetya B, Polii-Mandang J, Suryanto E. Phytochemical analysis of eceng gondok (Eichhornia crassipes) as agricultural waste as biosensitizer for ferri photoreduction. *AGRIVITA Journal of Agricultural Science.* 2012;**34**(2). doi: [10.17503/AgriVita-2012-34-2-p152-160](https://doi.org/10.17503/AgriVita-2012-34-2-p152-160).
36. Tosun M, Ercisli S, Sengul M, Ozer H, Polat T, Ozturk E. Antioxidant properties and total phenolic content of eight Salvia species from Turkey. *Biol Res.* 2009;**42**(2):175-81. [PubMed: [19746262](https://pubmed.ncbi.nlm.nih.gov/19746262/)].
37. Aboul-Enein AM, Shanab SM, Shalaby EA, Zahran MM, Lightfoot DA, El-Shemy HA. Cytotoxic and antioxidant properties of active principals isolated from water hyacinth against four cancer cells lines. *BMC Complement Altern Med.* 2014;**14**:397. doi: [10.1186/1472-6882-14-397](https://doi.org/10.1186/1472-6882-14-397). [PubMed: [25315352](https://pubmed.ncbi.nlm.nih.gov/25315352/)]. [PubMed Central: [PMC4213547](https://pubmed.ncbi.nlm.nih.gov/PMC4213547/)].
38. Badhani B, Sharma N, Kakkar R. Gallic acid: a versatile antioxidant with promising therapeutic and industrial applications. *RSC Advances.* 2015;**5**(35):27540-57. doi: [10.1039/c5ra01911g](https://doi.org/10.1039/c5ra01911g).
39. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci.* 2004;**74**(17):2157-84. doi: [10.1016/j.lfs.2003.09.047](https://doi.org/10.1016/j.lfs.2003.09.047). [PubMed: [14969719](https://pubmed.ncbi.nlm.nih.gov/14969719/)]. [PubMed Central: [PMC7126989](https://pubmed.ncbi.nlm.nih.gov/PMC7126989/)].
40. Milde J, Elstner EF, Grassmann J. Synergistic inhibition of low-density lipoprotein oxidation by rutin, gamma-terpinene, and ascorbic acid. *Phytomedicine.* 2004;**11**(2-3):105-13. doi: [10.1078/0944-7113-00380](https://doi.org/10.1078/0944-7113-00380). [PubMed: [15070159](https://pubmed.ncbi.nlm.nih.gov/15070159/)].
41. Lalitha P, Sripathi SK, Jayanthi P. Secondary metabolites of Eichhornia crassipes (Waterhyacinth): A review (1949 to 2011). *Nat Prod Commun.*

- 2012;7(9):1249–56. [PubMed: [23074922](#)].
42. Surendraraj A, Farvin KHS, Anandan R. Antioxidant potential of water hyacinth (*Eichhornia crassipes*): In vitro antioxidant activity and phenolic composition. *J Aquat Food Prod Technol*. 2013;22(1):11–26. doi: [10.1080/10498850.2011.621582](#).
 43. Labarrere B, Prinzing A, Dorey T, Chesneau E, Hennion F. Variations of secondary metabolites among natural populations of subantarctic ranunculus species suggest functional redundancy and versatility. *Plants (Basel)*. 2019;8(7). doi: [10.3390/plants8070234](#). [PubMed: [31331007](#)]. [PubMed Central: [PMC6681328](#)].
 44. Bobbarala V. Antifungal activity of selected plant extracts against phytopathogenic fungi *Aspergillus niger* F2723. *Indian J Sci Technol*. 2009;2(4):87–90. doi: [10.17485/ijst/2009/v2i4.15](#).
 45. Fareed MF, Haroon AM, Rabeh SA. Antimicrobial activity of some macrophytes from Lake Manzalah (Egypt). *Pak J Biol Sci*. 2008;11(21):2454–63. doi: [10.3923/pjbs.2008.2454.2463](#). [PubMed: [19205264](#)].
 46. Zhou B, Peng JP, Guo JW, Tang SB. Research on the antibacterial activities of extract from *Eichhornia crassipes*. *Jiangsu Journal of Agricultural Sciences*. 2009;25(3):547–50.
 47. Bouarab Chibane L, Degraeve P, Ferhout H, Bouajila J, Oulahal N. Plant antimicrobial polyphenols as potential natural food preservatives. *J Sci Food Agric*. 2019;99(4):1457–74. doi: [10.1002/jsfa.9357](#). [PubMed: [30206947](#)].
 48. Daglia M. Polyphenols as antimicrobial agents. *Curr Opin Biotechnol*. 2012;23(2):174–81. doi: [10.1016/j.copbio.2011.08.007](#). [PubMed: [21925860](#)].
 49. Cushnie TP, Lamb AJ. Recent advances in understanding the antibacterial properties of flavonoids. *Int J Antimicrob Agents*. 2011;38(2):99–107. doi: [10.1016/j.ijantimicag.2011.02.014](#). [PubMed: [21514796](#)].