



Transfer of Learning from a Virtual Table Tennis Game to Drinking from a Glass Task in Stroke Patients

Jacilda Oliveira Passos ¹, Deyvson Paiva Brito¹, Debora Carvalho Oliveira ¹, Lorena Raquel Dantas de Macedo Borges¹, Pedro Passos ², Aline Braga Galvao Silveira Fernandes ³ and Tania Fernandes Campos ^{1,*}

¹Departamento de Fisioterapia, Universidade Federal do Rio Grande do Norte, Natal/RN, Brasil

²CIPER, Faculdade de Motricidade Humana, University of Lisbon, Portugal

³Faculdade de Ciências da Saúde do Trairi, Universidade Federal do Rio Grande do Norte, Santa Cruz/RN, Brasil

*Corresponding author: Departamento de Fisioterapia, Universidade Federal do Rio Grande do Norte, Lagoa Nova, Natal, RN, Brasil. Email: campostf@gmail.com

Received 2020 June 28; Revised 2021 March 06; Accepted 2021 March 08.

Abstract

Background: Given that stroke is the cause of several neurological impairments, it is necessary to evaluate interventions that can improve patients' motor performance and functionality.

Objectives: This study aimed to evaluate the motor learning transfer of a training program using a virtual table tennis game. We hypothesized that playing the virtual game may improve the performance of stroke patients in a functional activity (drinking from a glass). Moreover, this virtual reality training may have different effects according to the brain injury side.

Methods: A non-randomized controlled study was performed. Sampling was by convenience. Evaluation and reevaluation of a drinking from a glass task were conducted through kinematic analysis (transfer test). The training was carried out with XBOX 360 table tennis game (20 attempts). Performance (number of hits) in the game and average angular velocity of the drinking task were analyzed by ANOVA.

Results: The study included 20 hemiparesis patients (10 left), mean age 50.6 (SD 9.2) years, and 20 healthy individuals, mean age 50.9 (SD 8.8) years. Patients showed worse performance than healthy subjects; however, they improved performance between the first and last attempts ($P = 0.001$). In the transfer test, left hemiparetic patients exhibited significant improvement in the shoulder ($P = 0.0001$) and elbow average angular velocity ($P = 0.0001$), but this did not occur in right hemiparetic patients (shoulder: $P = 0.761$; elbow: $P = 0.666$).

Conclusions: The results suggest that the virtual game has clinical potential in the motor learning transfer process to a functional activity in the real world, mainly for patients with right brain injury.

Keywords: Stroke, Upper Extremity, Virtual Reality Exposure Therapy, Psychomotor Performance

1. Background

Stroke-related motor impairments do not allow the patient to accurately perform coordinated movements, which significantly affects upper limb use in basic activities of daily living (ADL) (e.g., eating, drinking, dressing, grooming) (1). Impaired movement velocity is an important cause of disability, since it is related to the neuromuscular reaction triggered by the central nervous system in response to environmental demands (2).

Stroke survivors have different impairments in the affected upper limb according to the laterality of the brain injury. Left hemisphere damage causes greater impairment in movement trajectory, while right hemisphere

damage has more impact on the extent of movement (3).

Virtual reality (VR) games have been used for upper limb rehabilitation in stroke patients to promote multi-sensory, motor, and cognitive stimulation through an interactive and motivational environment (4). Training in a virtual environment aims to transfer the skills acquired there to the real world (5). The transfer of improved function to ADL performance is an important aspect to consider in post-stroke rehabilitation (6).

The transfer of learning effects in VR to real environments is inconclusive, with less favorable results for non-immersive systems (users experience the virtual environment as observers and interact with it using devices that do not fully encompass sensory perceptions, resulting in a

lesser sense of immersion in the virtual world) and more favorable results for systems with a higher level of immersion (users interact more with the environment and the computer-generated stimuli and providing visual, auditory, and haptic sensation, enabling individuals to feel immersed in the virtual environment) (6, 7). The laterality of the injury was investigated to determine its influence on the motor learning process in virtual reality training for upper limb rehabilitation of stroke patients (8). However, this functional hemispheric specialization was not studied in the transfer of learning from the virtual to the real world.

2. Objectives

In this context, the aim of this study was to evaluate the motor learning transfer of a training program using a virtual table tennis game. Our hypothesis was that playing a virtual tennis game may improve the performance of stroke patients in an upper limb functional activity (drinking from a glass). Moreover, this VR training may have different effects according to the brain injury side.

3. Methods

3.1. Study Design and Ethic Aspects

This is a non-randomized controlled study which was approved by the Research Ethics Committee of the Federal University of Rio Grande do Norte (CAAE: 40436514.0.0000.5292), considering the ethical principles of the Declaration of Helsinki. All participants were informed of the research procedures and provided written informed consent to participate in the study.

3.2. Participants

The sample size was calculated using the comparison of two averages of the angular velocity of the elbow in the reaching movement during the water-drinking activity of stroke patients ($64.9^\circ/\text{s}$) and healthy individuals ($121.8^\circ/\text{s}$), obtained from the Alt Murphy study (9). With these averages, we used a spreadsheet based on Singer's calculations (10). We found $n=10$ for each group, considering a 95% confidence level and 99% statistical power.

Sampling was by convenience. Individuals who were enrolled in the physiotherapy outpatient service were evaluated and selected according to the inclusion criteria. Healthy individuals were university employees recruited

and matched for sex, age, and education based on the characteristics of each patient included in the study until the sample was complete.

The study consisted of ten left (LP) and ten right (RP) hemiparetic patients. Ten healthy individuals who underwent left upper limb (LUL) and ten right upper limb (RUL) training were assessed.

Eligibility criteria for sample inclusion were: (i) first-episode stroke diagnosis; (ii) unilateral brain injury; (iii) chronic stage of stroke (after 6 months); (iv) right-handed individuals; (v) aged ≤ 70 years; (vi) able to respond to clinical assessment instruments and (vii) move the affected upper limb to perform the drinking task, necessary for kinematic analysis. Those with signs of unilateral neglect, evaluated by the National Institute of Health Stroke Scale validated by Lyden (11), and those with cognitive impairment assessed by the Mini-Mental State Examination who scored below the cutoff point established by Brucki et al (12) for the Brazilian population were excluded from the study.

3.3. Procedures

The patients were clinically assessed using the following instruments: NIHSS to evaluate the degree of neurological impairment, with scores ranging from 0 to 42 and the following cutoff points: 0 to 5 = light; 6 to 13 = moderate; 14 or more = severe; the Functional Independence Measure to assess items such as self-care, sphincter control, mobility and locomotion used in the study conducted by Kimura et al (13); the upper limb sub-item of the Fugl-Meyer scale to evaluate motor function, developed and validated by Fugl-Meyer (14); and the modified Ashworth scale to assess muscle tone in the shoulder and elbow, according to Biering-Sørensen et al (15).

After the evaluation, subjects trained on the XBOX 360 Kinect device (Microsoft®, Redmond, Washington, USA). Patients underwent training with the paretic upper limb. To maintain the pairing of healthy individuals, they trained with the right or left upper limb, according to the corresponding paired patient. The Kinect Sports® "paddle panic" mini game was used for training. In this game, the participant's avatar bats 60 balls that another player throws with only one hand in 45 seconds. Every hit ball was computed as one point. To hit the ball the individual performs shoulder flexion with forward and sideways elbow extension, depending on the direction of the ball. Patients performed 2 sets of 10 trials each. The rest period was 45 seconds between each trial and 15 minutes between sets, totaling 30 minutes of training (Figure 1). Healthy individuals performed only 1 trial of the game (60 balls in 45

seconds) in order to compare with the evolution of the patients' performance (Figure 1).

The average angular velocity of the shoulder and elbow in the sagittal plane (flexion and extension) during the task of drinking from a glass, was obtained through kinematic analysis, before and after the virtual game (Figure 1). The task was recorded using the Qualisys Motion Capture System (16) by capturing (12) reflective markers positioned on the upper limb, according to Qualisys Track Manager 2.6 - QTM guide. The markers were named and exported to Visual 3D software, where a biomechanical model was created, and the processing of the kinematic analysis was conducted.

To perform the proposed activity, participants remained seated on a 46cm-high bench and their feet were fully supported on the floor. In front of the participant, a table 73cm-high was placed. The table was at a distance where the hand remained supported on the edge of the table, the elbow in approximately 90° of flexion and the arm in a vertical position and close to the body. This distance allowed participants to perform the activity without rotating or bending their torso. A 200mL acrylic cup filled with 100mL of water was placed in the center of the table. The drinking task involved reaching for the glass, picking it up, lifting it off the table, drinking some water, returning it to the center of the table and placing their hand in the initial position.

3.4. Statistical analysis

The statistical package SPSS 22.0 (Statistical Package for the Social Science) was used for statistical analysis, assigning a significance level of 5%. The Shapiro-Wilk test was used to verify the distribution of data. The unpaired Student's *t*-test was used to confirm the pairing of healthy individuals and patients and compare data from the clinical evaluation of patients with right and left hemiparesis. Repeated-measures ANOVA with the Bonferroni post-hoc test were used to determine performance differences between trials in the virtual table tennis game and determine differences in average angular velocity in the drinking task between patients and healthy individuals, before and after training (evaluation and reevaluation). Cohen's *d* was calculated to verify effect size, according to the following classification: ($> = 0.15$ and < 0.40); ($> = 0.40$ and < 0.75); ($> = 0.75$ and < 1.10); ($> = 1.10$ and < 1.45), that is, small, medium, large, very large and huge effect, respectively (17).

4. Results

According to data analysis, there was no significant difference in age, weight, height, and education between patients and healthy individuals (Table 1). LP and LUL groups were composed of 5 men and 5 women. RP and RUL groups consisted of 3 men and 7 women. There was no significant difference between LP and RP groups in time since stroke, NIHSS (neurological impairment), FIM (functional independence), Fugl-Meyer scale (upper limb motor impairment) and Ashworth scores (spasticity) (Table 1). Patients exhibited mild neurological impairment and moderate motor impairment in the affected upper limb.

In regard to performance in the virtual table tennis game, a significant difference was observed between the groups of healthy individuals and patients on the first attempt (LUL x LP; $P = 0.0001$) (RUL x RP; $P = 0.001$), but not the last (LUL x LP; $P = 0.716$) (RUL x RP; $P = 0.723$). The patients showed a significant difference between the first and last attempt in the virtual table tennis game (LP: $P = 0.0001$ and RP: $P = 0.001$) (Figure 2).

Kinematic analysis indicated a significant difference between the evaluation of the LUL and LP groups in average shoulder angular velocity and at reevaluation ($P = 0.0001$) (Table 2). There was a significant difference between the evaluation of the LP group and at reevaluation, with an increase in angular velocity after training ($P = 0.0001$) (Table 2). In relation to elbow angular velocity, analysis revealed a significant difference between the LUL and LP groups evaluation ($P = 0.0001$) and at reevaluation ($P = 0.0001$). There was a significant difference between LP group evaluation and reevaluation ($P = 0.028$).

With respect to of the RUL and RP groups, a significant difference was observed in average shoulder angular velocity at evaluation ($P = 0.0001$) and reevaluation ($P = 0.0001$); however, there was no difference between RP group evaluation and reevaluation ($P = 0.761$). For the elbow, a difference in average angular velocity was noted between RUL and RP evaluation ($P = 0.0001$) and reevaluation ($P = 0.0001$). There was also no significant difference for the elbow between RP evaluation and reevaluation ($P = 0.666$).

Effect size analysis was only performed when significant intergroup differences were found. The effect size decreased in LUL and LP evaluation and reevaluation for both the shoulder and elbow. On the other hand, the effect size of the RUL and RP groups increased between evaluation and reevaluation of the shoulder and decreased for the elbow (Table 3).




Figure 1. Flowchart of the Methodological Procedures

Table 1. Comparison of Demographic and Anthropometric Data Between Patients (LP - Left Hemiparesis and RP - Right Hemiparesis) and Healthy Groups (LUL - Training with the Left Upper Limb) (RUL - Training with the Right Upper Limb) and Comparison of Clinical Data of the Patients. Data Presented by Mean and Standard Deviation (SD).

	LP (N = 10)	LUL (N = 10)	P-Value	RP (N = 10)	RUL (N = 10)	P-Value
Age (y)	52.2 (9.7)	54.6 (8.1)	0.56	49.8 (7.7)	47.7 (9.2)	0.11
Education (y)	10.8 (4.0)	11.9 (2.2)	0.34	9.2 (2.5)	11.2 (1.0)	0.37
Weight (kg)	77.3 (11.7)	70.7 (9.6)	0.11	68.3 (10.6)	62.4 (12.6)	0.15
Height (cm)	1.6 (0.8)	1.6 (0.8)	0.42	1.6 (0.6)	1.6 (0.7)	0.53
Post-stroke time (months)	24.6 (11.4)			25.1 (14.1)		0.93
NIHSS (score)	2.4 (1.1)			3.1 (2.1)		0.37
Motor FIM (score)	72.6 (6.4)			73.5 (4.8)		0.72
Fugl-Meyer (score)	49.9 (9.7)			50.8 (7.1)		0.81
Ashworth (score)						
Shoulder	1.7 (0.5)			1.5 (0.8)		0.53
Elbow	1.8 (0.8)			2.0 (0.8)		0.58

Abbreviations: NIHSS: National Institute Health Stroke Scale; FIM: Functional Independence Measure.

5. Discussion

The study aimed to evaluate the transfer of learning from a virtual table tennis game to the functional activity of drinking from a glass. Considering hemispheric specialization, we also aimed to determine if there are differ-

ences between patients with left and right brain injury in terms of transfer of learning. The main result obtained was that, after training, LP improved their performance in the transfer activity when compared to the initial assessment, achieving transfer of learning from a virtual game




Figure 2. Comparison of the performance (number of hits) between healthy individuals who underwent training with left (LUL) and right upper limb (RUL) and patients with left (LP) and right hemiparesis (RP), in first and last trials of virtual table tennis game.

Table 2. Comparison of Shoulder and Elbow Average Angular Velocity in Evaluation of Healthy Individuals Who Underwent Training with Their Left Upper Limb (LUL) and Healthy Individuals Who Trained Their Right Upper Limb (RUL) and in Evaluation and Reevaluation of Patients with Left (LP) and Right Hemiparesis (RP). Data Presented by Mean and Standard Deviation (SD).

	LUL Evaluation	LP Evaluation	LP Reevaluation
Average angular velocity (°/s)			
Shoulder	9.0 (3.8) ^{a,b}	4.9 (2.7) ^c	6.3 (3.2)
Elbow	18.9 (4.2) ^{a,b}	11.6 (3.9) ^c	13.6 (4.4)
	RUL Evaluation	RP Evaluation	RP Reevaluation
Average angular velocity (°/s)			
Shoulder	8.5 (2.4) ^{a,b}	7.0 (4.2)	6.4 (1.7)
Elbow	20.4 (9.1) ^{a,b}	12.9 (4.0)	13.7 (2.3)

^a Significant difference between the evaluation of the healthy individuals and evaluation of the hemiparesis patients.

^b Significant difference between the evaluation of the healthy individuals and reevaluation of the hemiparesis patients.

^c Significant difference between the evaluation and reevaluation of the hemiparesis patients.

to a functional activity, but RP did not.

Although previous studies have highlighted the influence of virtual environment fidelity on the transfer from virtual to real environments (7), in the present study, the functional activity that was the basis for the transfer test was not the one that was trained (table tennis). This leads us to believe that there may be a transfer of learning from the real to the virtual environment for functional

activities different from those trained in a virtual environment. Moreover, hemispheric specialization may have influenced this process.

Accordingly, virtual reality games can promote greater activation of the right brain hemisphere, especially in areas of somatosensory association, due to the multisensory stimulation that they provide with visuospatial feedback (18). These findings may explain the less significant im-

Table 3. Effect Size by D' Cohens for Average Angular Velocity of Shoulder and Elbow in the Studied Groups

	D' Cohens	Effect Size
Shoulder		
LUL evaluation x LP evaluation	1.30	very large
LUL evaluation x LP reevaluation	0.82	large
LP evaluation x LP reevaluation	0.48	medium
Elbow		
LUL evaluation x LP evaluation	1.89	huge
LUL evaluation x LP reevaluation	1.24	very large
LP evaluation x LP reevaluation	0.51	medium
Shoulder		
RUL evaluation x RP evaluation	0.48	medium
RUL evaluation x RP reevaluation	1.07	large
RP evaluation x RP reevaluation	-	-
Elbow		
RUL evaluation x RP evaluation	1.13	very large
RUL evaluation x RP reevaluation	0.98	large
RP evaluation x RP reevaluation	-	-

provement in average angular velocity after VR games in RP, corroborating other results showing that virtual games may have a better effect on LP (8).

In addition to the results already discussed, all the patients performed the table tennis virtual game worse than healthy subjects, clearly showing the influence of neurological and motor impairment on game performance. However, patient performance improved consistently, indicating a motor learning effect and activation of the central nervous system of these patients in favor of neural reorganization after training for a specific task with virtual table tennis (19).

This may have been caused by the varied practice that the game provided, since the launching of the table tennis balls was repetitive and random. The literature shows that this type of practice may result in the individual's performing better on the task (20). In this respect, varied practice must be considered during motor rehabilitation, because performance during the attempts may indicate whether the patient has evolved from the cognitive stage to associative and autonomous motor learning.

Due to their brain injury, the movement of stroke patients is expected to be slow, because of space-time decoupling and abnormal muscle activation patterns, making movements more segmented and variable than in healthy individuals (21). The process of learning transfer between motor skills is important for the therapist during clinical

practice (7), since it promotes an increase in the number of functional activities that can be performed at home and in the social life of patients (22). In addition, when rehabilitation involves the transfer of learning, it is able to integrate the "Activity" and "Participation" domains of the International Classification of Functionality, Disability and Health (ICF) (22). The ICF-based clinical practice must include promoting an individual's activity within ecological contexts integrating perception and action, and promoting a focus on the performer, task and environment (23).

The main limitation of the study was the short training time, which did not characterize the long-term effects of the learning transfer process. Future prospective studies should analyze the motor response of stroke patients submitted to a virtual reality protocol with longer training time and determine the long-term retention of motor skills, in order to better support clinical practices and taking into account the brain injury side.

5.1. Conclusion

According to our analysis, patients with right brain injury (left hemiparesis) improved shoulder and elbow angular velocity in the drinking from a glass task after training with a virtual table tennis game. However, patients with left brain injury (right hemiparesis) did not achieve this outcome. The results suggest that the virtual game has clinical potential in the motor learning transfer process to

a functional activity in the real world, mainly for patients with right brain injury.

Footnotes

Authors' Contribution: TFC and ABGSF participated of the conception and design of the study. JOP, DPB, DCO and LRDMB did the acquisition of data, analysis and interpretation of data. PP reviewed and approved the final version of the article to be submitted.

Conflict of Interests: The authors declare that they do not have any conflict of interest.

Ethical Approval: This is a non-randomized controlled study which was approved by the Research Ethics Committee of the Federal University of Rio Grande do Norte (CAAE: 40436514.0.0000.5292), considering the ethical principles of the Declaration of Helsinki. <http://plataformabrasil.saude.gov.br/login.jsf>

Funding/Support: This study was supported in part by Coodenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 and by National Council for Scientific and Technological Development CNPq (Grant number: 475182/2012-0).

Informed Consent: All participants were informed of the research procedures and provided written informed consent to participate in the study.

References

1. Wondergem R, Pisters MF, Wouters EJ, Olthof N, de Bie RA, Visser-Meily JM, et al. The course of activities in daily living: Who is at risk for decline after first ever stroke? *Cerebrovasc Dis*. 2017;**43**(1-2):1-8. doi: [10.1159/000451034](https://doi.org/10.1159/000451034). [PubMed: [27750246](https://pubmed.ncbi.nlm.nih.gov/27750246/)].
2. DeJong SL, Schaefer SY, Lang CE. Need for speed: Better movement quality during faster task performance after stroke. *Neurorehabil Neural Repair*. 2012;**26**(4):362-73. doi: [10.1177/1545968311425926](https://doi.org/10.1177/1545968311425926). [PubMed: [22140198](https://pubmed.ncbi.nlm.nih.gov/22140198/)]. [PubMed Central: [PMC3325333](https://pubmed.ncbi.nlm.nih.gov/PMC3325333/)].
3. Mani S, Mutha PK, Przybyla A, Haaland KY, Good DC, Sainburg RL. Contralesional motor deficits after unilateral stroke reflect hemisphere-specific control mechanisms. *Brain*. 2013;**136**(Pt 4):1288-303. doi: [10.1093/brain/aws283](https://doi.org/10.1093/brain/aws283). [PubMed: [23358602](https://pubmed.ncbi.nlm.nih.gov/23358602/)]. [PubMed Central: [PMC3613707](https://pubmed.ncbi.nlm.nih.gov/PMC3613707/)].
4. Perez-Marcos D. Virtual reality experiences, embodiment, videogames and their dimensions in neurorehabilitation. *J Neuroeng Rehabil*. 2018;**15**(1):113. doi: [10.1186/s12984-018-0461-0](https://doi.org/10.1186/s12984-018-0461-0). [PubMed: [30477527](https://pubmed.ncbi.nlm.nih.gov/30477527/)]. [PubMed Central: [PMC6258149](https://pubmed.ncbi.nlm.nih.gov/PMC6258149/)].
5. Nemani A, Ahn W, Cooper C, Schwaizberg S, De S. Convergent validation and transfer of learning studies of a virtual reality-based pattern cutting simulator. *Surg Endosc*. 2018;**32**(3):1265-72. doi: [10.1007/s00464-017-5802-8](https://doi.org/10.1007/s00464-017-5802-8). [PubMed: [28812196](https://pubmed.ncbi.nlm.nih.gov/28812196/)]. [PubMed Central: [PMC5809233](https://pubmed.ncbi.nlm.nih.gov/PMC5809233/)].
6. Kim WS, Cho S, Ku J, Kim Y, Lee K, Hwang HJ, et al. Clinical application of virtual reality for upper limb motor rehabilitation in stroke: Review of technologies and clinical evidence. *J Clin Med*. 2020;**9**(10). doi: [10.3390/jcm9103369](https://doi.org/10.3390/jcm9103369). [PubMed: [33096678](https://pubmed.ncbi.nlm.nih.gov/33096678/)]. [PubMed Central: [PMC7590210](https://pubmed.ncbi.nlm.nih.gov/PMC7590210/)].
7. Levac DE, Huber ME, Sternad D. Learning and transfer of complex motor skills in virtual reality: a perspective review. *J Neuroeng Rehabil*. 2019;**16**(1):121. doi: [10.1186/s12984-019-0587-8](https://doi.org/10.1186/s12984-019-0587-8). [PubMed: [31627755](https://pubmed.ncbi.nlm.nih.gov/31627755/)]. [PubMed Central: [PMC6798491](https://pubmed.ncbi.nlm.nih.gov/PMC6798491/)].
8. Fernandes A, de Melo JCP, de Oliveira DC, Cavalcanti F, Postolache OA, Passos PJM, et al. Is motor learning of stroke patients in non-immersive virtual environment influenced by laterality of injury? A preliminary study. *J Bodyw Mov Ther*. 2021;**25**:53-60. doi: [10.1016/j.jbmt.2020.10.015](https://doi.org/10.1016/j.jbmt.2020.10.015). [PubMed: [33714511](https://pubmed.ncbi.nlm.nih.gov/33714511/)].
9. Alt Murphy M, Willen C, Sunnerhagen KS. Kinematic variables quantifying upper-extremity performance after stroke during reaching and drinking from a glass. *Neurorehabil Neural Repair*. 2011;**25**(1):71-80. doi: [10.1177/1545968310370748](https://doi.org/10.1177/1545968310370748). [PubMed: [20829411](https://pubmed.ncbi.nlm.nih.gov/20829411/)].
10. Singer J. A simple procedure to compute the sample size needed to compare two independent groups when the population variances are unequal. *Stat Med*. 2001;**20**(7):1089-95. doi: [10.1002/sim.722](https://doi.org/10.1002/sim.722). [PubMed: [11276038](https://pubmed.ncbi.nlm.nih.gov/11276038/)].
11. Lyden P. Using the National Institutes of Health Stroke Scale: A Cautionary Tale. *Stroke*. 2017;**48**(2):513-9. doi: [10.1161/STROKEAHA.116.015434](https://doi.org/10.1161/STROKEAHA.116.015434). [PubMed: [28077454](https://pubmed.ncbi.nlm.nih.gov/28077454/)].
12. Brucki SM, Nitrini R, Caramelli P, Bertolucci PH, Okamoto IH. [Suggestions for utilization of the mini-mental state examination in Brazil]. *Arq Neuropsiquiatr*. 2003;**61**(3B):777-81. doi: [10.1590/s0004-282x2003000500014](https://doi.org/10.1590/s0004-282x2003000500014). [PubMed: [14595482](https://pubmed.ncbi.nlm.nih.gov/14595482/)].
13. Kimura Y, Yamada M, Hamanaka K, Tanaka N, Muroh Y. Usefulness of the prediction method based on a logarithmic model for functional recovery in stroke patients: in case of using the motor-Functional Independence Measure score. *Int J Rehabil Res*. 2017;**40**(2):134-7. doi: [10.1097/MRR.0000000000000219](https://doi.org/10.1097/MRR.0000000000000219). [PubMed: [28177943](https://pubmed.ncbi.nlm.nih.gov/28177943/)].
14. Fugl-Meyer AR, Jaasko L, Leyman I, Olsson S, Steglind S. The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance. *Scand J Rehabil Med*. 1975;**7**(1):13-31. [PubMed: [1135616](https://pubmed.ncbi.nlm.nih.gov/1135616/)].
15. Biering-Sorensen F, Nielsen JB, Klinge K. Spasticity-assessment: a review. *Spinal Cord*. 2006;**44**(12):708-22. doi: [10.1038/sj.sc.3101928](https://doi.org/10.1038/sj.sc.3101928). [PubMed: [16636687](https://pubmed.ncbi.nlm.nih.gov/16636687/)].
16. Qualisys Medical AB. *Qualisys Track Manager 2.6*. Sweden: Gothenburg; 2011. Available from: <https://www.qualisys.com/software/qualisys-track-manager/>.
17. Thalheimer W, Cook S. How to calculate effect sizes from published research: A simplified methodology. *Work-Learning Research*. 2009:1-9.
18. Oliveira SMS, Medeiros CSP, Pacheco TBF, Bessa N, Silva FGM, Tavares NSA, et al. Electroencephalographic changes using virtual reality program: technical note. *Neurol Res*. 2018;**40**(3):160-5. doi: [10.1080/01616412.2017.1420584](https://doi.org/10.1080/01616412.2017.1420584). [PubMed: [29291703](https://pubmed.ncbi.nlm.nih.gov/29291703/)].
19. Dabrowski J, Czajka A, Zielinska-Turek J, Jaroszynski J, Furtak-Niczyporuk M, Mela A, et al. Brain Functional Reserve in the Context of Neuroplasticity after Stroke. *Neural Plast*. 2019;**2019**:9708905. doi: [10.1155/2019/9708905](https://doi.org/10.1155/2019/9708905). [PubMed: [30936915](https://pubmed.ncbi.nlm.nih.gov/30936915/)]. [PubMed Central: [PMC6415310](https://pubmed.ncbi.nlm.nih.gov/PMC6415310/)].
20. Moliterno AH, Bezerra FV, Pires LA, Roncolato SS, da Silva TD, Massetti T, et al. Effect of Contextual Interference in the Practicing of a Computer Task in Individuals Poststroke. *Biomed Res Int*. 2020;**2020**:2937285. doi: [10.1155/2020/2937285](https://doi.org/10.1155/2020/2937285). [PubMed: [32775414](https://pubmed.ncbi.nlm.nih.gov/32775414/)]. [PubMed Central: [PMC7396124](https://pubmed.ncbi.nlm.nih.gov/PMC7396124/)].
21. Trombly CA. Deficits of reaching in subjects with left hemiparesis: a pilot study. *Am J Occup Ther*. 1992;**46**(10):887-97. doi: [10.5014/ajot.46.10.887](https://doi.org/10.5014/ajot.46.10.887). [PubMed: [1463060](https://pubmed.ncbi.nlm.nih.gov/1463060/)].
22. Lexell J, Brogardh C. The use of ICF in the neurorehabilitation process. *NeuroRehabilitation*. 2015;**36**(1):5-9. doi: [10.3233/NRE-141184](https://doi.org/10.3233/NRE-141184). [PubMed: [25547759](https://pubmed.ncbi.nlm.nih.gov/25547759/)].

23. Vaz DV, Silva PL, Mancini MC, Carello C, Kinsella-Shaw J. Towards an ecologically grounded functional practice in rehabilitation. *Hum Mov Sci.* 2017;**52**:117–32. doi: [10.1016/j.humov.2017.01.010](https://doi.org/10.1016/j.humov.2017.01.010). [PubMed: [28187353](https://pubmed.ncbi.nlm.nih.gov/28187353/)].