Published online 2021 August 29.

Research Article



Height Prediction Based on the Lengths of Ulna and Tibia in an Iranian Population

Shiva Mokhtari¹, Mahmoud Hajiahmadi ¹, Haleh Esmaili ¹, and Reza Ghadimi ¹, ³, ^{*}

¹Student Research Committee, Babol University of Medical Sciences, Babol, Iran

²Non-Communicable Pediatric Diseases Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

³Social Determinants of Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

corresponding author: Social Determinants of Health Research Center, Health Research Center, Babol University of Medical Sciences, Ganjafrooz St., Babol, Postal Code: 47176-47754, Iran. Email: rezaghadimi@yahoo.com

Received 2020 August 28; Revised 2021 May 23; Accepted 2021 May 23.

Abstract

Background: Height is an important factor for medical, nutrition, and forensic sciences; nevertheless, measuring proves to be a challenging task in some cases. In this respect, an alternative immediate, accurate, and possible anthropometric evaluation is needed.

Objectives: This study was set to find a proper formula to estimate height from the lengths of the ulna and tibia in Iranian adults. **Methods:** A total of 500 healthy males and females aged 20-40 years were randomly selected from the volunteers' pool for this cross-sectional study. Ulna and tibia lengths and standing heights were measured according to standard protocols. Ulna and tibia lengths were applied to find a reliable equation to predict stature accurately. Data were statistically analyzed by SPSS version 17 using regression, curve estimation, and linear model.

Results: The mean (SD) heights of male and female participants were 176.45 (11.98) cm and 161.29 (10.11) cm, respectively, while the right and left ulna and the right and left tibia were 29.05 (1.63) cm, 29.03 (1.44) cm, 38.86 (1.33) cm, and 38.88 (1.25) cm, respectively. The correlation coefficients of $r^2 = 0.80$ and $r^2 = 0.69$ for males and females participants' right ulna, respectively, showed a significant correlation with height. Hence, the new formula provided reliable results for stature estimation for northern Iran subgroups. **Conclusions:** Equations based based on right ulna length are more reliable and accurate for height prediction in both genders. It should be considered that these equations could be different among ethnically diverse populations, even in the northern Iranian population.

Keywords: Stature Prediction, Body Height, Ulna, Tibia, Anthropometry

1. Background

Stature as a physical characteristic is one of the principal parameters of identification and assessment (1, 2). Height is a fundamental health indicator especially used to assess the medicinal requirements of hospitalized patients. However, in some cases, including people with particular disorders, people of age, and hospitalized patients, the concept of height becomes an intricate challenge for having a direct and accurate assessment (3, 4). The relationship between height and different body parts such as the head, trunk, and upper and lower limbs has always been an interesting subject for anatomists and anthropologists; moreover, it can also be very useful for forensic medicine experts, especially in providing them with a statistical database for anthropometrical measurements in future studies (2, 5-7). For this purpose, various screening tools have been introduced and applied, complementing

other variables, including medical, clinical, biochemical, anthropometric, and dietary measurements (8). Regardless of the availability of these tools, the accuracy of height measurement when acquired with certain difficulties becomes inherently questionable (9, 10). Hence, it is essential to devise alternative methods to measure stature accurately and easily. In this manner, several studies have explored a variety of methods such as using arm span, knee height, and ulna length to find a reliable formula that gives an approximate estimate for stature when the actual height is difficult to be measured (11-15). Furthermore, height as a central anthropometric measure can be influenced by various factors, such as genetics, ethnicity, and other environmental circumstances (16, 17). In this respect, it is necessary to find an applicable and suitable equation to screen patients from each nation or ethnic group or to identify the absence of this equation among people of dif-

Copyright © 2021, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

ferent parts of each region like the northern population of Iran. Since ulna and tibia lengths can be easily measured, and considering the availability and inexpensiveness of related tools, the method appears as a suitable alternative for height estimation.

2. Objectives

This study aimed to offer a proper formula for the prediction of height based on the lengths of the ulna and tibia of adults in north of Iran.

3. Methods

3.1. Study Population

This study was approved by the Medical Research Ethics Committee of Babol University of Medical Sciences (No: 9441667). The subjects were selected from a pool of 20 to 40-year-old staff, students, and patients at Babol Health center. People who were left-handed and those with any kinds of bone surgery, fractures, physical disabilities, and skeletal malformations that could potentially influence height or ulna and tibia measurements were excluded. Overall, 528 subjects were selected using the simple random sampling method from the admission list. Twentyeight subjects were excluded based on the exclusion criteria. The final sample of 500 subjects was divided into two groups based on gender (250 males versus 250 females). In line with the code of conduct, the study was explained to all the participants, and afterwards, consent forms were signed by them prior to the start of the study.

3.2. Anthropometric Measurements

All assessments were performed between 9.00 a.m. and 4 p.m. with the same equipment and trained technicians to minimize methodological errors and potential biases. Height was measured by a portable stadiometer (Iran Teb, Tehran) that was mounted on the wall. Height was measured for each participant to the nearest 0.1 cm while standing in erect position, facing forward, with Frankfort horizontal head position and bare feet without a hat or a hairstyle that could affect the process of measurement (18). The forearm (ulna) was measured using an anthropometric tape (Sewing meter, China) from the point of olecranon extending to the point of the wrist (styloid). For each participant, the left and right forearms were measured while flexed across the chest and in front of the body without any objects like a wristband, a bracelet, or a watch (13). The measurements were repeated three times and were eventually recorded to the nearest 0.1 cm. Left and right tibia lengths were measured by a sewing meter and were recorded to

the nearest 0.1 cm. It was considered as the distance between the upper border of the medial condyle and the tip of the left medial malleolus, while the knee was kept at a 90-degree angle in a sitting position.

3.3. Statistical Analysis

Data analysis was performed using SPSS version 18. The mean measurement was reported as mean plus standard deviation. Pearson's correlation coefficient and simple linear regression were run to examine the relationship between ulna and tibia lengths and height and stature in both genders. A P-value of less than 0.05 was considered significant.

4. Results

The mean (SD) heights of the male and female participants were 176.45 (11.9) cm and 161.29 (10.11) cm, respectively. In men, the mean (SD) lengths of the right and left ulna were 29.05 (1.63) cm and 29.03 (1.44) cm, respectively, and they were 27.74 (1.49) cm and 27.78 (1.45) cm in women, respectively. Moreover, the mean lengths of the right and left tibia were respectively 38.86 (1.33) cm and 38.88 (1.25) cm in men and 35.89 (1.01) and 35.88 (1.09) in women.

Table 1 shows the relationship between the actual height and ulna and tibia lengths. There was a significant difference between the mean of ulna and tibia in confidence level of adjusted R square for select the equation. Height formula was suggested for right ulna length in both genders as it shows the highest level of adjusted R square with an impressive significance level (P < 0.01). Moreover, ulna and tibia in men and women were significantly different level (Table 1).

The correlation between right ulna length and height was moderately strong and significant (r = 0.833 for women and r = 0.897 for men in the right ulna). Therefore, the height estimation formulae for men and women were considered as follows:

Predicted height [cm] = $88.71 + (3.02 \times \text{ulna length} \text{[cm]})$ and Predicted height [cm] = $46.41 + (4.28 \times \text{ulna length} \text{[cm]})$, respectively.

Although there was a significant relationship between the length of the tibia and height, the relationship was somewhat weak in both genders ($r^2 = 0.66 \& r^2 = 0.56$ in women and men, respectively, P < 0.001). Other equations were also reliable, as presented in Table 1.

Figures 1 and 2 display the scatter diagram related to ulna and tibia lengths from both left and right sides against height for each gender. The solid line represents the least-squares best-fit regression and height estimation based on the related regression equation. The majority

Table 1. Correlation Between the Measured Height and Suggested Equations for Predicted Height Based on Ulna and Tibia Lengths by Gender							
Group	Correlation Between Height with Ulna and Tibia		Regression Coefficients			Predicted Height (cm)	F
	Adj. r ²	P-value	A Slope Constant	В	95%CI	— Mean (SD)	Equation
Male							
Right ulna	0.805	< 0.001	3.02	88.71	2.83 - 3.20	176.0 (4.3)	Predicted height [cm] = 88.70 + (3.02 × ulna length [cm])
Left ulna	0.792	< 0.001	2.96	90.58	2.76 - 3.14	175.6 (5.8)	Predicted height [cm] = 90.57+(2.95 × ulna length [cm])
Right tibia	0.568	< 0.001	3.54	38.93	3.25 - 3.80	179.1 (5.4)	Predicted height [cm] = 38.93 + (3.53 × ulna length [cm])
Left tibia	0.561	< 0.001	3.47	41.22	3.27 - 3.61	180.7 (4.8)	Predicted height [cm] = 41.22 + (3.74 × ulna length [cm])
Female							
Right ulna	0.693	< 0.001	4.28	46.41	4.04 - 4.41	166.3 (1.9)	Predicted height [cm] = 46.41+(4.27 × ulna length [cm])
Left ulna	0.671	< 0.001	4.07	51.98	3.88 - 4.43	164.1 (3.2)	Predicted height [cm] = 51.98+(4.06 × ulna length [cm])
Right tibia	0.667	< 0.001	2.04	87.91	1.76 - 2.34	168.4 (4.0)	Predicted height [cm] = 87.91 + (2.04 × ulna length [cm])
Left tibia	0.667	< 0.001	2.04	88.05	1.82 - 2.36	161.0 (3.80)	Predicted height [cm] = $88.05+(2.03 \times ulna$ length [cm])

Abbreviations: Adj. r², Adjusted R square; SD, standard deviation; CI, confidence intervals.

of the values for the right ulna subgroups in both groups were above the predicted regression. The overall trend of the right ulna agreed more with the equation prediction in both genders, such that these concordances were supported by the regression coefficients (Table 1).

Height prediction using the four variables (right and left ulna and tibia lengths) by multivariate regression analysis showed (adj. $r^2 = 0.804$ and $r^2 = 0.831$ in women and men, respectively) the almost similar correlation coefficients for the stature estimation based on right ulna length.

5. Discussion

This study was carried out to examine the appropriate equations for stature estimation and prediction using ulna and tibia lengths in healthy Iranian adults. The results showed that the correlation coefficients between height and ulna length were higher than those the tibia length in both genders. Moreover, between right and left ulna

lengths, right ulna length with a greater correlation coefficient proved to be more accurate for height prediction in men and women in the north of Iran. In this regard, some other studies were performed to investigate reliable stature formulae according to long bone length. It had been understood that stature equations differ based on various factors such as gender, ethnic group, and study methods (19-22). Madden et al. (2012) claimed that their formula for assessing the height based on the length of ulna correlated closely with the actual height in white participants, while it had failed to provide an accurate assessment of the actual height in black and Asian participants (4). Moreover, according to other studies, it has been revealed that the stature formulae of people can vary, especially considering those who live in different parts of the same state (20, 23). In this study, the participants were of similar ethnicity and were residing in the same region. Moreover, there is also the theory that a person's lifestyle, namely urban or rural life, childhood nutrition and growth, and type of job, can affect stature. According to the available



Figure 1. Distribution of the measured height against ulna (a) and tibia (b) lengths in men. The solid line represents the regression for the values from the participants.



Figure 2. Distribution of the measured height against ulna (a) and tibia (b) lengths in women. The solid line represents the regression for the values from the participants.

evidence, several environmental factors, such as vigorous physical activity, infectious diseases, hormonal disorders, malnutrition, and poor living conditions, can prevent the individual from reaching the genetically determined adult height, especially caused by a decrease in the growth of legs (24). Hence, stature estimation and prediction based on the upper limbs may not be reliable among those populations. Therefore, all conditions should be considered to provide the appropriate formula to estimate height.

Since easy and fast equations for the calculation of stature are key in malnutrition screening, examining a bedridden patient, and forensic identifications, multiplevariable equations by multivariate regression analysis were not performed. In this study, left-handed subjects were not included due to the availability of right-handed people, and the unavailability of an acceptable left-handed sample size within the time of research. Although there was not a difference between the lengths of the right and the left ulna, there is a concern about the use of these equations, especially considering different lengths of ulna between the dominant and the non-dominant hands in lefthanded people, which affects equations and the accuracy of prediction. In this regard, our results may not be useful for left-handed individuals considering the dominant hand as a significant factor in the measurement of ulna lengh. We, therefore, highly recommend another study to be carried out among left-handed people.

5.1. Conclusions

It could be concluded that the equations for predicting height based on ulna length, especially the right ulna in adults, provides a reliable estimation of height in healthy adults located in the north of Iran.

Footnotes

Authors' Contribution: Study concept and design: Shiva Mokhtari and Reza Ghadimi; Data acquisition: Reza Ghadimi and Shiva Mokhtari; Analysis and interpretation of the data: Reza Ghadimi, Shiva Mokhtari, and Haleh Esmaili; Drafting of the manuscript: Shiva Mokhtari and Haleh Esmaili; Critical revision of the manuscript for important intellectual content: Haleh Esmaili and Reza Ghadimi; Statistical analysis: Mahmoud Hajiahmadi; Administrative, technical, and material support: Mahmoud Hajiahmadi and Reza Ghadimi; Study supervision: Mahmoud Hajiahmadi and Reza Ghadimi.

Conflict of Interests: We have no conflicts of interest to disclose.

Ethical Approval: This study was approved by the Medical Research Ethics Committee of Babol University of Medical Sciences (Code NO: 9441667).

Funding/Support: The present research received financial support from the Research Deputy of Babol University of Medical Sciences.

Informed Consent: Consent forms were signed by participants prior to the initiation of the study.

References

- Menendez Garmendia A, Sanchez-Mejorada G, Gomez-Valdes JA. Stature estimation formulae for Mexican contemporary population: A sample based study of long bones. J Forensic Leg Med. 2018;54:87–90. doi: 10.1016/j.jflm.2017.12.019. [PubMed: 29331714].
- 2. Kaore DA, Kaore DB, Kamdi DA, kaore DS. Stature estimation from tibial length: Stature estimation from tibial length. *Natl J Integr Res Med.* 2012;**3**(2):45–50.
- Ilayperuma I, Nanayakkara G, Palahepitiya N. A model for the estimation of personal stature from the length of forearm. *Int J Morphol.* 2010;28(4):1081–6. doi: 10.4067/s0717-9502201000400015.
- Madden AM, Tsikoura T, Stott DJ. The estimation of body height from ulna length in healthy adults from different ethnic groups. J Hum Nutr Diet. 2012;25(2):121–8. doi: 10.1111/j.1365-277X.2011.01217.x. [PubMed: 22077418].
- Gonzalez-Colmenares G, Medina CS, Baez LC. Estimation of stature by cephalometric facial dimensions in skeletonized bodies: study from a sample modern Colombians skeletal remains. *Forensic Sci Int.* 2016;**258**:101 e1–6. doi: 10.1016/j.forsciint.2015.10.016. [PubMed: 26631845].
- Gualdi-Russo E, Bramanti B, Rinaldo N. Stature estimation from tibia percutaneous length: New equations derived from a Mediterranean population. *Sci Justice*. 2018;**58**(6):441–6. doi: 10.1016/j.scijus.2018.08.001. [PubMed: 30446073].
- Rodriguez S, Miguens X, Rodriguez-Calvo MS, Febrero-Bande M, Munoz-Barus JI. Estimating adult stature from radiographically determined metatarsal length in a Spanish population. *Forensic Sci Int.* 2013;**226**(1-3):297 e1-4. doi: 10.1016/j.forsciint.2012.12.006. [PubMed: 23312585].
- Green SM, Watson R. Nutritional screening and assessment tools for use by nurses: literature review. J Adv Nurs. 2005;50(1):69–83. doi: 10.1111/j.1365-2648.2004.03350.x. [PubMed: 15788067].
- Cook Z, Kirk S, Lawrenson S, Sandford S. Use of BMI in the assessment of undernutrition in older subjects: reflecting on practice. *Proc Nutr* Soc. 2005;64(3):313-7. doi: 10.1079/pns2005437. [PubMed: 16048662].
- Kirk SFL, Hawke T, Sandford S, Wilks Z, Lawrenson S. Are the measures used to calculate BMI accurate and valid for the use in older people? *Journal of Human Nutrition and Dietetics*. 2003;16(5):366–7. doi: 10.1046/j.1365-277X.2003.04673.x.
- Agnihotri AK, Kachhwaha S, Jowaheer V, Singh AP. Estimating stature from percutaneous length of tibia and ulna in Indo-Mauritian population. *Forensic Sci Int.* 2009;**187**(1-3):109 e1-3. doi: 10.1016/j.forsciint.2009.02.010. [PubMed: 19286337].
- Brown JK, Whittemore KT, Knapp TR. Is arm span an accurate measure of height in young and middle-age adults? *Clin Nurs Res*. 2000;**9**(1):84– 94. doi: 10.1177/10547730022158456. [PubMed: 11271050].
- Gauld LM, Kappers J, Carlin JB, Robertson CF. Height prediction from ulna length. *Dev Med Child Neurol.* 2004;46(7):475–80. doi: 10.1017/s0012162204000787.
- Hickson M, Frost G. A comparison of three methods for estimating height in the acutely ill elderly population. *J Hum Nutr Diet.* 2003;16(1):13-20. doi: 10.1046/j.1365-277x.2003.00416.x. [PubMed: 12581405].
- Ritz P. Validity of measuring knee-height as an estimate of height in diseased French elderly persons. *J Nutr Health Aging*. 2004;8(5):386–8. [PubMed: 15359357].

- Chumlea WC, Guo SS, Wholihan K, Cockram D, Kuczmarski RJ, Johnson CL. Stature prediction equations for elderly non-Hispanic white, non-Hispanic black, and Mexican-American persons developed from NHANES III data. J Am Diet Assoc. 1998;98(2):137–42. doi: 10.1016/S0002-8223(98)00036-4. [PubMed: 12515412].
- Launer LJ, Harris T. Weight, height and body mass index distributions in geographically and ethnically diverse samples of older persons. Ad Hoc Committee on the Statistics of Anthropometry and Aging. *Age Ageing*. 1996;**25**(4):300–6. doi: 10.1093/ageing/25.4.300. [PubMed: 8831875].
- Samson MM, Meeuwsen IB, Crowe A, Dessens JA, Duursma SA, Verhaar HJ. Relationships between physical performance measures, age, height and body weight in healthy adults. *Age Ageing*. 2000;29(3):235–42. doi: 10.1093/ageing/29.3.235. [PubMed: 10855906].
- 19. Borhani-Haghighi M, Navid S, Hassanzadeh G. Height prediction from ulnar length in Chabahar: A city in South-East of Iran. *Romanian Journal of Legal Medicine*. 2016;**24**(4):304–7. doi: 10.4323/rjlm.2016.304.
- 20. Ghanbaril K, Nazari AR, Ghanbari A, Chehrei S. Stature estimation and

formulation of based on ulna length in Kurdish racial subgroup. *Ital J Anat Embryol.* 2016;**121**(1):43–50. [PubMed: 28872796].

- 21. Navid S, Mokhtari T, Alizamir T, Arabkheradmand A, Hassanzadeh G. [Determination of stature from upper arm length in medical students]. *Anat Sci J.* 2014;**11**(3):135–40. Persian.
- Saco-Ledo G, Porta J, Duyar I, Mateos A. Stature estimation based on tibial length in different stature groups of Spanish males. *Forensic Sci Int.* 2019;**304**:109973. doi: 10.1016/j.forsciint.2019.109973. [PubMed: 31605880].
- 23. Gupta P, Kumar P, Gaharwar A, Ansari H, Hussein M. Correlation of percutaneous length of tibia with body height and estimation of stature in living North Indian males. *Sch. J. App. Med. Sci.* 2014;**2**(2D):848–52.
- Bogin B, Varela-Silva MI. Leg length, body proportion, and health: a review with a note on beauty. *Int J Environ Res Public Health*. 2010;7(3):1047-75. doi: 10.3390/ijerph7031047. [PubMed: 20617018]. [PubMed Central: PMC2872302].